Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.

aSF601
.A1U5
c2

United States Department of Agriculture

Animal and Plant Health Inspection Service

Veterinary Services

APHIS 91-40
May 1982

National Poultry Improvement Plan and Auxiliary Provisions

USDA
NAT'L AGRIC. LIBRARY
MAY 5 '88



NATIONAL POULTRY IMPROVEMENT PLAN

The National Poultry Improvement Plan became operative July 1, 1935, with the approval of the Secretary of Agriculture and under the authority of an appropriation made by Congress for the United States Department of Agriculture to be used in cooperation with State authorities in the administration of regulations for the improvement of poultry, poultry products, and hatcheries. The National Turkey Improvement Plan became operative September 25, 1943. Based on the recommendations of the 1970 National Plans Conference, the two Plans were consolidated into one National Poultry Improvement Plan with general provisions applicable to all classes of domesticated fowl and with special provisions applicable to problems and conditions peculiar to particular classes of such fowl. Authority for the administration of the Plan is now contained in the Department of Agriculture Organic Act of 1944, as amended (7 U.S.C. 429).

The objective of the National Poultry Improvement Plan is to provide a cooperative State-Federal program through which new technology can be effectively applied to the improvement of poultry and poultry products throughout the country. The provisions of the Plan, developed jointly by industry members and State and Federal officials, establish standards for the evaluation of poultry breeding stock and hatchery products with respect to freedom from hatchery-disseminated diseases. Products conforming to specific standards are identified by authorized terms that are uniformly applicable in all parts of the country.

The provisions of the Plan are changed from time to time to conform with the development of the industry and to utilize new information as it becomes available. These changes are based upon recommendations made at the National Plan Conferences by official delegates representing participating flockowners, breeders, and hatcherymen from all cooperating States, in accordance with Subpart E, Part 147 of this publication.

Acceptance of the Plan is optional with the States and individual members of the industry within the States. The Plan is administered in each State by an Official State Agency cooperating with the United States Department of Agriculture. A list of the Official State Agencies and other publications relating to the Plan are available upon request to the National Poultry Improvement Plan, APHIS-VS, Building 265, BARC-East, Beltsville, Maryland 20705.

ANIMAL AND PLANT HEALTH INSPECTION SERVICE
VETERINARY SERVICES

CHAPTER I - ANIMAL AND PLANT HEALTH INSPECTION SERVICE
VETERINARY SERVICES[1/]

DEPARTMENT OF AGRICULTURE

SUBCHAPTER F - POULTRY IMPROVEMENT

Part
145 National Poultry Improvement Plan
146 [Reserved]
147 Auxiliary Provisions on National Poultry Improvement Plan

SUBCHAPTER F - POULTRY IMPROVEMENT

PART 145 - NATIONAL POULTRY IMPROVEMENT PLAN

Subpart A - General Provisions

Sec.
145.1 Definitions.
145.2 Administration.
145.3 Participation.
145.4 General provisions for all participants.
145.5 Specific provisions for participating flocks.
145.6 Specific provisions for participating hatcheries.
145.7 Specific provisions for participating dealers.
145.8 Terminology and classification; general.
Sec.
145.9 Terminology and classification; hatcheries and dealers.
145.10 Terminology and classification; flocks, products, and States.
145.11 Supervision.
145.12 Inspections.
145.13 Debarment from participation.
145.14 Blood testing.

Subpart B - Special Provisions for Egg-Type Chicken Breeding Flocks and Products

145.21 Definitions.
145.22 Participation.
145.23 Terminology and classification; flocks and products.
145.24 Terminology and classification; States.

1/ 44 Federal Register 61586, October 26, 1979.

Subpart C - Special Provisions for Meat-Type Chicken Breeding Flocks and Products

145.31 Definitions.
145.32 Participation.
145.33 Terminology and classification; flocks and products.
145.34 Terminology and classification; States.

Subpart D - Special Provisions for Turkey Breeding Flocks and Products

Sec.
145.41 Definitions.
145.42 Participation.
145.43 Terminology and classification; flocks and products.
145.44 Terminology and classification; States.

Subpart E - Special Provisions for Waterfowl, Exhibition Poultry, and Game Bird Breeding Flocks and Products

145.51 Definitions.
145.52 Participation.
145.53 Terminology and classification; flocks and products.
145.54 Terminology and classification; States.

AUTHORITY: Section 101(b) of the Department of Agriculture Organic Act of 1944, as amended, 58 Stat. 734 (7 U.S.C. 429)

Subpart A - General Provisions

§145.1 Definitions.

Words used in this part in the singular form shall be deemed to import the plural, and vice versa, as the case may demand. Except where the context otherwise requires, for the purposes of this part the following terms shall be construed, respectively, to mean:

(a) Plan. The provisions of the National Poultry Improvement Plan contained in this part.

(b) Person. A natural person, firm, or corporation.

(c) Department. The U.S. Department of Agriculture.

(d) Service. The Animal and Plant Health Inspection Service, Veterinary Services, of the Department.

(e) State. Any State, the District of Columbia, or Puerto Rico.

(f) Official State Agency. The State authority recognized by the Department to cooperate in the administration of the Plan.

(g) State Inspector. Any person employed or authorized under §145.11(b) to perform functions under this part.

(h) Authorized Agent. Any person designated under §145.11(a) to perform functions under this part.

(i) Affiliated flockowner. A flockowner who is participating in the Plan through an agreement with a participating hatchery.

(j) Flock (1) As applied to breeding. All poultry of one kind of mating (breed and variety or combination of stocks) and of one classification on one form;

(j)(2) As applied to disease control. All of the poultry on one farm except that, at the discretion of the Official State Agency, any group of poultry which is segregated from another group and has been so segregated for a period of at least 21 days may be considered as a separate flock.

(k) Hatchery. Hatchery equipment on one premises operated or controlled by any person for the production of baby poultry.

(l) Poultry. Domesticated fowl, including chickens, turkeys, waterfowl, and game birds, except doves and pigeons, which are bred for the primary purpose of producing eggs or meat.

(m) Domesticated. Propagated and maintained under the control of a person.

(n) Products. Poultry breeding stock and hatching eggs, baby poultry, and started poultry.

(o) Baby poultry. Newly hatched poultry (chicks, poults, ducklings, goslings, keets, etc.) that have not been fed or watered.

(p) Started poultry. Young poultry (chicks, pullets, cockerels, capons, poults, ducklings, goslings, keets, etc.) that have been fed and watered and are less than 6 months of age.

(q) Strain. Poultry breeding stock bearing a given name produced by a breeder through at least five generations of closed flock breeding.

(r) Stock. A term used to identify the progeny of a specific breeding combination within a species of poultry. These breeding combinations may include pure strains, strain crosses, breed crosses, or combinations thereof.

(s) Primary breeding flock. A flock composed of one or more generations that is maintained for the purpose of establishing, continuing, or improving parent lines.

(t) Multiplier breeding flock. A flock that is intended for the production of hatching eggs used for the purpose of producing progeny for commercial egg or meat production or for other nonbreeding purposes.

(u) Trade name or number. A name or number compatible with State and Federal laws and regulations applied to a specified stock or product thereof.

(v) Franchise breeder. A breeder who normally sells products under a specific strain or trade name and who authorizes other hatcheries to produce and sell products under this same strain or trade name.

(w) Franchise hatchery. A hatchery which has been authorized by a franchise breeder to produce and sell products under the breeder's strain or trade name.

(x) Pullorum disease or pullorum. A disease of poultry caused by Salmonella pullorum.

(y) Fowl typhoid or typhoid. A disease of poultry caused by Salmonella gallinarum.

(z) S. typhimurium infection or typhimurium. A disease of poultry caused by Salmonella typhimurium or S. typhimurium var. copenhagen.

(aa) Official supervision. (1) As applied to Plan programs. The direction, inspection, and critical evaluation by the Official State Agency of compliance with the provisions of the Plan;

(2) As applied to non-Plan but equivalent State poultry improvement programs. The direction, inspection, and critical evaluation by an officer or agency of a State government, of compliance with a publicly announced State poultry improvement program.

(bb) Authorized laboratory. A laboratory designated by an Official State Agency, subject to review by the Service, to perform the blood testing and bacteriological examinations provided for in this part.

(cc) Salmonella. Any of the species of the bacteria belonging to the Salmonella genus, except that members of the arizona group are not included in this definition.

(dd) Colon bacilli. For the purpose of this chapter, those organisms which are gram negative, non spore-forming bacilli, which ferment lactose with gas formation, and serve as an index of fecal contamination.

(ee) Equivalent or equivalent requirements. Requirements which are equal to the program, conditions, criteria, or classifications with which compared, as determined by the Official State Agency and with the concurrence of the Service.

(ff) Sexual maturity. The average age at which a species of poultry is biologically capable of reproduction.

§145.2 Administration.

(a) The Department cooperates through a Memorandum of Understanding with Official State Agencies in the administration of the Plan.

(b) The administrative procedures and decisions of the Official State Agency are subject to review by the Service. The Official State Agency shall carry out the administration of the Plan within the State according to the applicable provisions of the Plan and the Memorandum of Understanding.

(c) An Official State Agency may accept for participation an affiliated flock located in another State under a mutual understanding and agreement, in writing, between the two Official State Agencies regarding conditions of participation and supervision.

(d) The Official State Agency of any State may, except as limited by §145.3(d), adopt regulations applicable to the administration of the Plan in such State further defining the provisions of the Plan or establishing higher standards compatible with the Plan.

§145.3 Participation.

(a) Any person producing or dealing in products may participate in the Plan when he has demonstrated, to the satisfaction of the Official State Agency, that his facilities, personnel, and practices are adequate for carrying out the applicable provisions of the Plan, and has signed an agreement with the Official State Agency to comply with the general and the applicable specific provisions of the Plan and any regulations of the Official State Agency under §145.2. Affiliated flockowners may participate without signing an agreement with the Official State Agency.

(b) Each participant shall comply with the Plan throughout the operating year of the Official State Agency, or until released by such Agency.

(c) A participant in any State shall participate with all of his poultry hatching egg supply flocks and hatchery operations within such State. He shall report to the Official State Agency on NPIP Form 3B or through other appropriate means each breeding flock before the birds reach 24 weeks of age. This report will include:

(c)(1) Name and address of flockowner;

(c)(2) Flock location and designation;

(c)(3) Type: Primary or Multiplier;

(c)(4) Breed, variety, strain, or trade name of stock;

(c)(5) Source of males;
(c)(6) Source of females;
(c)(7) Number of birds in the flock; and
(c)(8) Intended classification of flock.

(d) No person shall be compelled by the Official State Agency to qualify products for any of the other classifications described in §145.10 as a condition of qualification for the U.S. Pullorum-Typhoid Clean classification.

(e) Participation in the Plan shall entitle the participant to use the Plan emblem reproduced below:



Figure 1

§145.4 General provisions for all participants.

(a) Records of purchases and sales and the identity of products handled shall be maintained in a manner satisfactory to the Official State Agency.

(b) Products, records of sales and purchase of products, and material used to advertise products shall be subject to inspection by the Official State Agency at any time.

(c) Advertising must be in accordance with the Plan, and applicable rules and regulations of the Official State Agency and the Federal Trade Commission. A participant advertising products as being of any official classification may include in his advertising reference to associated or franchised hatcheries only when such hatcheries produce the same kind of products of the same classification.

(d) Participants may not buy, sell, or receive for any purpose products from nonparticipants, Provided, That with the permission of the Official State Agency and approval of the Service, other products may be bought or received for use in breeding flocks or for experimental purposes.

(e) Each participant shall be assigned a permanent approval number by the Service. This number, prefaced by the numerical code of the State, will be the official approval number of the participant and may be used on each certificate, invoice, shipping label, or other document used by the participant in the sale of his products. Each Official State Agency which requires an approval or permit number for out-of-State participants to ship into its State should honor this number. The approval number shall be withdrawn when the participant no longer qualifies for participation in the Plan.

§145.5 Specific provisions for participating flocks.

(a) Poultry equipment, and poultry houses and the land in the immediate vicinity thereof, shall be kept in sanitary condition as recommended in §§147.21 and 147.22(a) and (e) of this chapter. The participating flock, its eggs, and all equipment used in connection with the flock shall be separated from nonparticipating flocks, in a manner acceptable to the Official State Agency.

(b) All flocks shall consist of healthy, normal individuals characteristic of the breed, variety, cross, or other combination which they are stated to represent.

(c) A flock shall be deemed to be a participating flock at any time only if it has qualified for the U.S. Pullorum-Typhoid Clean classification, as prescribed in subpart B,C,D, or E of this part.

(d) Each bird shall be identified with a sealed and numbered band obtained through or approved by the Official State Agency: Provided, That exception may be made at the discretion of the Official State Agency.

§145.6 Specific provisions for participating hatcheries.

(a) Hatcheries, including brooder rooms, shall be kept in sanitary condition, acceptable to the Official State Agency. The procedures outlined in §§147.22 through 147.25 of this chapter shall be considered as a guide in determining compliance with this provision. The minimum requirements with respect to sanitation shall include the following:

(a)(1) Incubator walls, floors, and trays shall be kept free from broken eggs and eggshells.

(a)(2) Tops of incubators and hatchers shall be kept clean (not used for storage).

(a)(3) Entire hatchery, including sales room, shall be kept in a neat, orderly condition and free from accumulated dust.

(a)(4) Hatchery residue, such as eggshells, infertile eggs, and dead germs, shall be disposed of promptly and in a manner satisfactory to the Official State Agency.

(a)(5) Hatchers and hatching trays shall be cleaned and fumigated or disinfected after each hatch, preferably using the procedures outlined in §§147.24(b) and 147.25(e) of this chapter. While not mandatory for participation, all eggs set should be fumigated in accordance with the procedures recommended in §147.25 of this chapter.

(b) A hatchery which keeps started poultry must keep such poultry separated from the incubator room in a manner satisfactory to the Official State Agency.

(c) All baby and started poultry offered for sale under Plan terminology shall be normal and typical of the breed, variety, cross, or other combination represented.

(d) Eggs incubated shall be sound in shell, typical for the breed, variety, strain, or cross thereof and reasonably uniform in shape. Hatching eggs shall be trayed and the baby poultry boxed with a view to uniformity of size.

(e) All hatcheries within a State which are operated under the ownership or management of the same person or persons or related corporations shall participate in the Plan if any of them are to participate.

§145.7 Specific provisions for participating dealers.

Dealers in poultry breeding stock, hatching eggs, or baby or started poultry shall comply with all provisions in this part which apply to their operations.

§145.8 Terminology and classification; general.

(a) The official classification terms defined in §§145.9 and 145.10 and the various designs illustrative of the official classifications reproduced in §145.10 may be used only by participants and to describe products that have met all the specific requirements of such classifications.

(b) Products produced under the Plan shall lose their identity under Plan terminology when they are purchased for resale by or consigned to nonparticipants.

(c) Participating flocks, their eggs, and the baby and started poultry produced from them may be designated by their strain or trade name. When a breeder's trade name or strain designation is used, the participant shall be able by records to substantiate that the products so designated are from flocks that are composed of either birds hatched from eggs produced under the direct supervision of the breeder of such strain, or stock multiplied by persons designated and so reported by the breeder to each Official State Agency concerned.

§145.9 Terminology and classification; hatcheries and dealers.

Participating hatcheries and dealers shall be designated as "National Plan Hatchery" and "National Plan Dealer," respectively. All Official State Agencies shall be notified by the Service of additions, withdrawals, and changes in classification.

§145.10 Terminology and classification; flocks, products, and States.

Participating flocks, products produced from them, and States which have met the respective requirements specified in Part 145, Subpart B, C, D, or E may be designated by the following terms or illustrative designs:

(a) U.S. Approved. (See §145.53(a).)

Figure 2

(b) U.S. Pullorum-Typhoid Clean. (See §145.23(b), §145.33(b), §145.43(b), and §145.53(b).)

Figure 3

(c) U.S. M. Gallisepticum Clean. (See §145.23(c), §145.33(c), §145.43(c), and §145.53(c).)

Figure 4

(d) U.S. Sanitation Monitored. (See §145.23(d) and §145.33(d).)

Figure 5

(e) U.S. M. Synoviae Clean. (See §145.23(e) and §145.33(e).)

Figure 6

(f) U.S. M. Meleagridis Clean. (See §145.43(d).)

Figure 7

(g) U.S. Pullorum-Typhoid Clean State. (See §145.24(a), §145.34(a), §145.44(a), and §145.54(a).)



Figure 8

(h) U.S. Pullorum-Typhoid Clean State, Turkeys. (See §145.44(b).)

Figure 9

(i) U.S. Mycoplasma Gallisepticum Clean State, Turkeys. (See §145.44(c).)

Figure 10

§145.11 Supervision.

(a) The Official State Agency may designate qualified persons as Authorized Agents to do the sample collecting and blood testing provided for in §145.14 and the selecting required for the U.S. Approved classification provided for in §145.53(a).

(b) The Official State Agency shall employ or authorize qualified persons as State Inspectors to perform or supervise the performance of the selecting and testing of participating flocks, and to perform the official inspections necessary to verify compliance with the requirements of the Plan.

(c) Authorities issued under the provisions of this section shall be subject to cancellation by the Official State Agency on the grounds of incompetence or failure to comply with the provisions of the Plan or regulations of the Official State Agency. Such actions shall not be taken until a thorough investigation has been made by the Official State Agency and the authorized person has been given notice of the proposed action and the basis therefore and an opportunity to present his views.

§145.12 Inspections.

(a) Each participating hatchery shall be inspected a sufficient number of times each year to satisfy the Official State Agency that the operations of the hatchery are in compliance with the provisions of the Plan.

(b) Each year at least 15 percent of the independent flocks and the affiliated flocks of each hatchery shall be inspected by a State Inspector. Each inspection shall include the examination of a sufficient number of males and females and, in flocks qualified for participation by the whole-blood test, the blood testing of a sufficient number of birds to determine whether the work of the Authorized Agent was satisfactory and that the flock is qualified for participation. The State Inspector shall also determine whether or not the flock and premises are in compliance with the provisions in §145.5(a) and (b).

§145.13 Debarment from participation.

Participants in the Plan, who after investigation by the Official State Agency or its representative, are notified of their apparent noncompliance with the Plan provisions or regulations of the Official State Agency, shall be afforded a reasonable time, as specified by the Official State Agency, within which to demonstrate or achieve compliance. If compliance is not demonstrated

or achieved within the specified time, the Official State Agency may debar the participant from further participation in the Plan for such period, or indefinitely, as the Agency may deem appropriate. The debarred participant shall be afforded notice of the bases for the debarment and opportunity to present his views with respect to the debarment in accordance with procedures adopted by the Official State Agency. The Official State Agency shall thereupon decide whether the debarment order shall continue in effect. Such decision shall be final unless the debarred participant, within 30 days after the issuance of the debarment order, requests the Administrator to determine the eligibility of the debarred participant for participation in the Plan. In such event the Administrator shall determine the matter de novo in accordance with §§50.21 through 50.28-14 and §§50.30 through 50.33 of the rules of practice in 7 CFR Part 50, which are hereby made applicable to proceedings before the Administrator under this section. The definitions in 7 CFR 50.2(e), (g), (h), and (1) and the following definitions shall apply with respect to terms used in such rules of practice:

(a) "Administrator" means the Administrator, Animal and Plant Health Inspection Service of the U.S. Department of Agriculture or any officer or employee to whom authority has heretofore been delegated or to whom authority may hereafter be delegated to act in his stead.

§145.14 Blood testing.

Blood samples for official tests shall be drawn by an Authorized Agent or State Inspector and tested by an authorized laboratory, except that the stained-antigen, rapid whole-blood test for pullorum-typhoid may be conducted by an Authorized Agent or State Inspector. For those programs where a sample of the poultry may be tested in lieu of the entire flock, the poultry tested shall be a representative sample drawn from all pens or units of the flock.

(a) For Salmonella. (1) The official blood tests for pullorum-typhoid shall be the standard tube agglutination test, the microagglutination test, or the rapid serum test for all classes of poultry or the stained antigen, rapid whole-blood test for all classes of poultry except turkeys. The recommended procedures for conducting such tests are described in §§147.1, 147.2, 147.3, and 147.5 of this chapter. Each lot of antigen used for the whole-blood test shall be approved by the Department and shall be of the polyvalent type. All microtest antigens shall also be approved by the Department.

(a)(2) [Reserved]

(a)(3) There shall be an interval of at least 21 days between any official blood test and any previous test with salmonella antigen.

(a)(4) Poultry must be more than 4 months of age when tested: Provided, That candidates for participation under Subpart E of this part shall have attained the age of sexual maturity before being tested.

(a)(5) The official blood test shall include the testing of a sample of blood from each bird in the flock: Provided, That under specified conditions (see applicable provisions of §§145.23, 145.33, 145.43, and 145.53) the testing of a portion or sample of the birds may be used in lieu of testing each bird. When reactors are found in any flock, or S. pullorum or S. gallinarum isolations are made from baby poultry or fluff samples, the flock may qualify for participation with two consecutive official negative tests. Qualification of this flock, or any other flock on the same premises during the next 12 months, shall be based on the testing of all birds. Such testing shall be conducted by or directly supervised by a State Inspector.

(a)(6) All domesticated fowl, except waterfowl, on the farm of the participant shall either be properly tested to meet the same standards as the participating flock or these birds and their eggs shall be separated from the participating flock and its eggs.

(a)(7) All tests with salmonella antigens of flocks participating in or candidates for participation in the Plan shall be reported to the Official State Agency within 10 days following the completion of such tests. All reactors shall be considered in determining the classification of the flock.

(a)(8) Reactors shall be submitted to a laboratory for autopsy and bacteriological examination. All reactors in a flock if there are 4 or less reactors shall be submitted to a laboratory designated by the Official State Agency for bacteriological examination, as described in §147.11 of this chapter: Provided, That if more than 4 reactors are found, a minimum of 4 birds shall be submitted. The recommended minimum procedure for bacteriological examination is described in §147.11. When reactors are submitted within 10 days from date of reading the test and the bacteriological examination fails to demonstrate infection of the serotype for which the test was conducted, the flock shall be deemed to have had no reactors to the specified test.

(a)(9) Any drug, for which there is scientific evidence of masking the test reaction or hindering the bacteriological recovery of salmonella organisms, shall not be fed or administered to poultry within 3 weeks prior to a test or bacteriological examination upon which a salmonella classification is based.

(a)(10) When suitable evidence, as determined by the Official State Agency or the State Animal Disease Control Official, indicates that baby or started poultry produced by participating hatcheries are infected with organisms for which the parent flock received an official control classification and this evidence indicates that the infection was transmitted from the parent flock, the Official State Agency may, at its discretion, require additional testing of the flock involved. If infection is found in the parent flock, its classification shall be suspended until the flock is requalified under the requirements for the classification. Furthermore, the Official State Agency may require that the hatching eggs from such flocks be removed from the incubator and destroyed prior to hatching. When salmonella or arizona organisms are isolated from a specimen which originated in a participating hatchery, the Official State Agency shall attempt to locate the source of the infection. The results of the investigation and the action taken to eliminate the infection shall be reported by the Official State Agency to the Service.

(b) For M. gallisepticum and M. synoviae: (1) The official blood test for M. gallisepticum or M. synoviae shall be either the serum plate agglutination test, the tube agglutination test, the hemagglutination inhibition (HI) test, or a combination of two or more of these tests. The HI test shall be used to confirm the positive results of other serological tests. HI titers of 1:40 or less may be interpreted as equivocal, and final judgment may be based on further samplings and/or culture of reactors.

(b)(2) The tests shall be conducted using M. gallisepticum or M. synoviae antigens approved by the Department or the Official State Agency and shall be performed in accordance with the recommendations of the producer of the antigen.

(b)(3) When reactors to the test for which the flock was tested are submitted to a laboratory as prescribed by the Official State Agency, the criteria found in §147.6 shall be used in determining the final status of the flock.

(b)(4) Any drug, for which there is scientific evidence of masking the test reaction or hindering the bacteriological recovery of mycoplasma organisms, shall not be fed or administered to poultry within 3 weeks prior to a test or bacteriological examination upon which a mycoplasma classification is based.

Subpart B - Special Provisions For Egg-Type Chicken Breeding Flocks And Products

§145.21 Definitions.

Except where the context otherwise requires, for the purposes of this subpart the following terms shall be construed, respectively, to mean:

(a) Egg-type chicken breeding flocks. Flocks that are composed of stock that has been developed for egg production and are maintained for the principal purpose of producing chicks for the ultimate production of eggs for human consumption.

(b) Chicks. Newly hatched chickens which have not been fed or watered.

(c) Started chickens. Young chickens (chicks, pullets, cockerels, capons) which have been fed and watered and are less than 6 months of age.

§145.22 Participation.

Participating flocks of egg type chickens, and the eggs and chicks produced from them, shall comply with the applicable general provisions of Subpart A of this part and the special provisions of this Subpart B.

(a) The minimum weight of hatching eggs sold shall be 1 11/12 ounces each, except as otherwise specified by the purchaser of the eggs.

(b) Mediterranean breed eggs shall be reasonably free from tints.

(c) Started chickens shall lose their identity under Plan terminology when not maintained by Plan participants under the conditions prescribed in §145.5(a).

(d) Hatching eggs produced by primary breeding flocks shall be fumigated according to the procedures described in §147.25(a) of this chapter: Provided, That alternative sanitizing procedures may be used with the approval of the Official State Agency in each specific instance and with the general concurrence by the Service in the policy adopted by the Official State Agency.

§145.23 Terminology and classification; flocks and products.

Participating flocks, and the eggs and chicks produced from them, which have met the respective requirements specified in this section may be designated by the following terms and the corresponding designs illustrated in §145.10:

(a) [Reserved]

(b) U.S. Pullorum-Typhoid Clean. A flock in which freedom from pullorum and typhoid has been demonstrated to the Official State Agency under the criteria in one of the following paragraphs (b)(1) through (5) of this section: Provided, That a flock qualifying by means of a blood test shall be tested within the past 12 months, except that the retesting of a participating flock which is retained for more than 12 months shall be at the discretion of the Official State Agency with the concurrence of the Service. (See §145.14 relating to the official blood test where applicable.)

(b)(1) It has been officially blood tested with no reactors.

(b)(2) It is a multiplier breeding flock, or a breeding flock composed of progeny of a primary breeding flock which is intended solely for the production of multiplier breeding flocks, and meets the following specifications as determined by the Official State Agency and the Service:

(b)(2)(i) The flock is located in a State where all persons performing poultry disease diagnostic services within the State are required to report to the Official State Agency within 48 hours the source of all poultry specimens from which S. pullorum or S. gallinarum is isolated;

(b)(2)(ii) The flock is composed entirely of birds that originated from U.S. Pullorum-Typhoid Clean breeding flocks or from flocks that met equivalent requirements under official supervision; and

(b)(2)(iii) A sample comprised of at least 25 percent of the birds in the flock has been officially blood tested with no reactors, or its progeny has been subjected to an approved 10-day chick mortality bacteriological examination monitoring program and bacteriological examination of a sample of down shed by chicks in the hatcher from selected hatches as prescribed by the Official State Agency: Provided, That when the blood testing procedure is used, the percentage of the flock included in the sample may be reduced by 5 percentage points following each year in which there is no evidence of infection on the premises until the required percentage is reduced to zero: And provided further, That the sample tested for the qualification of a flock under this subparagraph shall include at least 500 birds the first year, 400 the second year, 300 the third year, 200 the fourth year, and 100 the fifth year.

(b)(3) It is a multiplier breeding flock, or a breeding flock composed or progeny of a primary breeding flock which is intended solely for the production of multiplier breeding flocks, that originated from U.S. Pullorum-Typhoid Clean breeding flocks or from flocks that met equivalent requirements under official supervision, and is located in a State in which it has been determined by the Service that:

(b)(3)(i) All hatcheries, except turkey hatcheries, within the State are qualified as "National Plan Hatcheries" or have met equivalent requirements for pullorum-typhoid control under official supervision;

(b)(3)(ii) All hatchery supply flocks, except turkey flocks, within the State, are qualified as U.S. Pullorum-Typhoid Clean or have met equivalent requirements for pullorum-typhoid control under official supervision: Provided, That if other domesticated fowl, except waterfowl, are maintained on the same premises as the participating flock, freedom from pullorum-typhoid infection shall be demonstrated by an official blood test of each of these fowl;

(b)(3)(iii) All shipments of products other than U.S. Pullorum-Typhoid Clean, or equivalent, into the State are prohibited;

(b)(3)(iv) All persons performing poultry disease diagnostic services within the State are required to report to the Official State Agency within 48 hours the source of all poultry specimens from which S. pullorum or S. gallinarum is isolated;

(b)(3)(v) All reports of S. pullorum or S. gallinarum isolations from poultry are promptly followed by an investigation by the Official State Agency to determine the origin of the infection;

(b)(3)(vi) All flocks found to be infected with pullorum or typhoid are quarantined until marketed or destroyed under the supervision of the Official State Agency, or until subsequently blood tested, following the procedure for reacting flocks as contained in §145.14(a)(5), and all birds fail to demonstrate pullorum or typhoid infection;

(b)(3)(vii) All poultry, including exhibition, exotic, and game birds, but excluding waterfowl, going to public exhibition shall come from U.S. Pullorum-Typhoid Clean or equivalent flocks, or have had a negative pullorum-typhoid test within 90 days of going to public exhibition;

(b)(3)(viii) Discontinuation of any of the condition or procedures described in paragraphs (b)(3)(i), (ii), (iii), (iv), (v), (vi), and (vii) of this section, or the occurrence of repeated outbreaks of pullorum or typhoid in poultry breeding flocks, other than turkey flocks, within or originating within the State shall be grounds for the Service to revoke its determination that such conditons and procedures have been met or complied with. Such action shall not be taken until a thorough investigation has been made by the Service and the Official State Agency has been given an opportunity to present its views.

(b)(4) It is a multiplier breeding flock located in a State which has been determined by the Service to be in compliance with the provisions of (b)(3) of this section, and in which pullorum disease or fowl typhoid is not known to exist nor to have existed in hatchery supply flocks, other than turkey, waterfowl, exhibition poultry, and game bird supply flocks, within the State during the preceding 12 months.

(b)(5) It is a primary breeding flock located in a State determined to be in compliance with the provisions of paragraph (b)(4) of this section, and in which a sample of 300 birds from flocks of more than 300, and each bird in flocks of 300 or less, has been officially tested for pullorum-typhoid with no reactors: Provided, That a bacteriological examination monitoring program acceptable to the Official State Agency and approved by the Service may be used in lieu of blood testing.

(c) U.S. M. Gallisepticum Clean. (1) A flock maintained in compliance with the provisions of §147.26 of this chapter and in which freedom from M. gallisepticum has been demonstrated under the criteria specified in paragraph (c)(1)(i) or (ii) of this section.

(c)(1)(i) It is a flock in which all birds or a sample of at least 300 birds has been tested for M. gallisepticum as provided in §145.14(b) when more than 4 months of age: Provided, That to retain this classification, a minimum of 150 birds shall be tested at intervals of not more than 90 days: And provided further, That a sample comprised of less than 150 birds may be tested at any one time, with the approval of the Official State Agency and the concurrence of the Service, provided that a minimum of 150 birds is tested within each 90-day period; or

(c)(1)(ii) It is a multiplier breeding flock which originated as U.S. M. Gallisepticum Clean chicks from primary breeding flocks and from which a sample comprised of a minimum of 150 birds per flock has been tested for M. gallisepticum as provided in §145.14(b) when more than 4 months of age: Provided, That to retain this classification, the flock shall be subjected to one of the following procedures:

(c)(1)(ii)(a) At intervals of not more than 90 days, a sample of 75 birds, with a minimum of 30 birds per pen, whichever is greater, shall be tested; or

(c)(1)(ii)(b) At intervals of not more than 30 days, a sample of 25 cull chicks produced from the flock shall be subjected to laboratory procedures acceptable to the Official State Agency and approved by the Service, for the detection and recovery of M. gallisepticum; or

(c)(2) A participant handling U.S. M. Gallisepticum Clean products shall keep these products separate from other products in a manner satisfactory to the Official State Agency: Provided, That U.S. M. Gallisepticum Clean chicks from primary breeding flocks shall be produced in incubators and hatchers in which only eggs from flocks qualified under paragraph (c)(1)(i) of this section are set.

(c)(3) U.S. M. Gallisepticum Clean chicks shall be boxed in clean boxes and delivered in trucks that have been cleaned and disinfected as described in §147.24(a) of this chapter.

(d) U.S. Sanitation Monitored. This program is intended to be the basis from which the breeding-hatching industry may conduct a program for the prevention and control of Salmonellosis. It is intended to reduce the incidence of Salmonella organisms in hatching eggs and chicks through an effective and practical sanitation program at the breeder farm and in the hatchery. This will afford other segments of the poultry industry an opportunity to reduce the incidence of Salmonella in their products.

(d)(1) A flock and the hatching eggs and chicks produced from it which have met the following requirements as determined by the Official State Agency:

(d)(1)(i) The flock shall originate from a source where sanitation and management practices, as outlined in §145.23(d)(1) of this paragraph, are conducted;

(d)(1)(ii) The flock is maintained in compliance with §§147.21, 147.24(a), and 147.26 of this chapter;

(d)(1)(iii) All feed should be pelletized and/or crumbled in mills operated at 190°F. or above in facilities for recontamination control;

(d)(1)(iv) Feed shall be stored and transported in such a manner as to prevent possible contamination;

(d)(1)(v) Hatching eggs are collected at least four times a day and are handled as described in §147.22 and fumigated on the farm as described in §147.25(a) of this chapter or handled in a sanitary manner equivalent to §§147.22 and 147.25(a) of this part, which has been approved by the Official State Agency with the concurrence of the Service; and

(d)(1)(vi) Chicks are hatched in a hatchery meeting the requirements of §§147.23 and 147.24(b) and are in compliance with §147.25(b) of this chapter.

(d)(2) The Official State Agency may use the procedures described in §147.14 of this chapter to monitor the effectiveness of the sanitation practices.

(d)(3) In order for a hatchery to sell products of this classification, all products handled shall meet the requirements of the classification.

(d)(4) This classification may be revoked by the Official State Agency if the participant fails to follow recommended corrective measures.

(e) U.S. M. Synoviae Clean. (1) A flock maintained in compliance with the provisions of §147.26 and in which freedom from M. synoviae has been demonstrated under the criteria specified in paragraph (e)(1)(i) or (ii) of this section.

(e)(1)(i) It is a flock in which a minimum of 300 birds has been tested for M. synoviae as provided in §145.14(b) when more than 4 months of age: Provided, That to retain this classification, a sample of at least 150 birds shall be tested at intervals of not more than 90 days: And provided further, That a sample comprised of less than 150 birds may be tested at any one time, with the approval of the Official State Agency and the concurrence of the Service, provided that a minimum of 150 birds is tested within each 90-day period; or

(e)(1)(ii) It is a multiplier breeding flock which originated as U.S. M Synoviae Clean chicks from primary breeding flocks and from which a sample comprised of a minimum of 75 birds has been tested for M. synoviae as provided in §145.14(b) when more than 4 months of age: Provided, That to retain this classification, a sample of 50 birds shall be tested at intervals or not more than 90 days: And provided further, That a sample of less than 50 birds may be tested at any one time, provided that a minimum of 30 birds per flock with a minimum of 15 birds per pen, whichever is greater, is tested each time and a total of at least 50 birds is tested within each 90-day period.

(e)(2) A participant handling U.S. M. Synoviae Clean products shall keep these products separate from other products in a manner satisfactory to the Official State Agency: Provided, That U.S. M. Synoviae Clean chicks from primary breeding flocks shall be produced in incubators and hatchers in which only eggs from flocks qualified under paragraph (e)(1)(i) or (ii) of this section are set.

(e)(3) U.S. M. Synoviae Clean chicks shall be boxed in clean boxes and delivered in trucks that have been cleaned and disinfected as described in §145.24(a).

(f) U.S. M. Gallisepticum Clean Started Poultry. (1) A flock which originated from U.S. M. Gallisepticum Clean breeding flocks and was hatched in a hatchery approved by the Official State Agency for the production of U.S. M. Gallisepticum Clean chicks.

(f)(2) All other poultry on the premises of the candidate flock must originate from U.S. M. Gallisepticum Clean sources.

(f)(3) The flock is maintained in compliance with the provisions of §147.26.

(f)(4) The flock's freedom from M. gallisepticum is demonstrated by a negative blood test, as provided in §145.14(b), of a sample of 75 birds, with a minimum of 50 birds per poultry house, between 15-20 days prior to the flock being moved to laying quarters.

(f)(5) Started poultry shall be delivered to and from the farm premises in crates and vehicles which have been cleaned and disinfected as described in §147.24(a) of this chapter.

§145.24 Terminology and classification; States.

(a) U.S. Pullorum-Typhoid Clean State. (1) A State will be declared a U.S. Pullorum-Typhoid Clean State when it has been determined by the Service that:

(a)(1)(i) The State is in compliance with the provisions contained in §145.23(b)(3)(i) through (vii), §145.33(b)(3) (i) through (vii), §145.43(b)(3)(i) through (vi), and §145.53(b)(3)(i) through (vii).

(a)(1)(ii) No pullorum disease or fowl typhoid is known to exist nor to have existed in hatchery supply flocks within the State during the preceding 12 months: Provided, That pullorum disease or fowl typhoid found in waterfowl, exhibition poultry, and game bird breeding flocks will not prevent a State, which is otherwise eligible, from qualifying for a period of 2 years.

(a)(2) Discontinuation of any of the conditions described in paragraph (a)(1)(i) of this section, or repeated outbreaks of pullorum or typhoid occur in hatchery supply flocks described in paragraph (a)(1)(ii) of this section, or if an infection spreads from the originating premises, the Service shall have grounds to revoke its determination that the State is entitled to this classification. Such action shall not be taken until a thorough investigation has been made by the Service and the Official State Agency has been given an opportunity for a hearing.

Subpart C - Special Provisions For Meat-Type Chicken Breeding Flocks And Products

§145.31 Definitions.

Except where the context otherwise requires, for the purposes of this subpart the following terms shall be construed, respectively, to mean:

(a) Meat-type chicken breeding flocks. Flocks that are composed of stock that has been developed for meat production and are maintained for the principal purpose of producing chicks for the ultimate production of meat.

(b) Chicks. Newly hatched chickens which have not been fed or watered.

(c) Started chickens. Young chickens (chicks, pullets, cockerels, capons) which have been fed and watered and are less than 6 months of age.

§145.32 Participation.

Participating flocks of meat-type chickens, and the eggs and chicks produced from them, shall comply with the applicable general provisions of Subpart A of this part and the special provisions of this Subpart C.

(a) The minimum weight of hatching eggs sold shall be 1 10/12 ounces each, except as otherwise specified by the purchaser of the eggs.

(b) Started chickens shall lose their identity under Plan terminology when not maintained by Plan participants under the conditions prescribed in §145.5(a).

(c) Hatching eggs produced by primary breeding flocks shall be fumigated according to the procedures described in §147.25(a): Provided, That alternative sanitizing procedures may be used with the approval of the Official State Agency in each specific instance and with the general concurrence by the Service in the policy adopted by the Official State Agency.

§145.33 Terminology and classification; flocks and products.

Participating flocks, and the eggs and chicks produced from them, which have met the respective requirements specified in this section may be designated by the following terms and the corresponding designs illustrated in §145.10:

(a) [Reserved]

(b) U.S. Pullorum-Typhoid Clean. A flock in which freedom from pullorum and typhoid has been demonstrated to the Official State Agency under the criteria in one of paragraphs (b)(1) through (5) of this section: Provided, That a flock qualifying by means of a blood test shall be tested within the past 12 months, except that the retesting of a participating flock which is retained for more than 12 months shall be at the discretion of the Official State Agency with the concurrence of the Service. (See §145.14 relating to the official blood test where applicable.)

(b)(1) It has been officially blood tested with no reactors.

(b)(2) It is a multiplier breeding flock, or a breeding flock composed of progeny of a primary breeding flock which is intended solely for the production of multiplier breeding flocks, and meets the following specifications as determined by the Official State Agency and the Service:

(b)(2)(i) The flock is located in a State where all persons performing poultry disease diagnostic services within the State are required to report to the Official State Agency within 48 hours the source of all poultry specimens from which S. pullorum or S. gallinarum is isolated;

(b)(2)(ii) The flock is composed entirely of birds that originated from U.S. Pullorum-Typhoid Clean breeding flocks or from flocks that met equivalent requirements under official supervison; and

(b)(2)(iii) A sample comprised of at least 25 percent of the birds in the flock has been officially blood tested with no reactors, or its progeny has been subjected to an approved 10-day chick mortality bacteriological examination monitoring program and bacteriological examination of a sample of down shed by chicks in the hatcher from selected hatches as prescribed by the Official State Agency: Provided, That when the blood testing procedures is used, the percentage of the flock included in the sample may be reduced by 5 percentage points following each year in which there is no evidence of infection on the premises until the required percentage is reduced to zero: And provided further, That the sample tested for the qualification of a flock under this subparagraph shall include at least 500 birds the first year, 400 the second year, 300 the third year, 200 the fourth year, and 100 the fifth year.

(b)(3) It is a multiplier breeding flock, or a breeding flock composed of progeny of a primary breeding flock which is intended solely for the production of multiplier breeding flocks, that originated from U.S. Pullorum-Typhoid Clean breeding flocks or from flocks that met equivalent requirements under official supervision, and is located in a State in which it has been determined by the Service that:

(b)(3)(i) All hatcheries, except turkey hatcheries, within the State are qualified as "National Plan Hatcheries" or have met equivalent requirements for pullorum-typhoid control under official supervision;

(b)(3)(ii) All hatchery supply flocks, except turkey flocks, within the State, are qualified as U.S. Pullorum-Typhoid Clean or have met equivalent requirements for pullorum-typhoid control under official supervision: Provided, That if other domesticated fowl, except waterfowl, are maintained on the same premises as the participating flock, freedom from pullorum-typhoid infection shall be demonstrated by an official blood test of each of these fowl;

(b)(3)(iii) All shipments of products other than U.S. Pullorum-Typhoid Clean, or equivalent, into the State are prohibited;

(b)(3)(iv) All persons performing poultry disease diagnostic services within the State are required to report to the Official State Agency within 48 hours the source of all poultry specimens from which S. pullorum or S. gallinarum is isolated;

(b)(3)(v) All reports of S. pullorum or S. gallinarum isolations from poultry are promptly followed by an investigation by the Official State Agency to determine the origin of the infection;

(b)(3)(vi) All flocks found to be infected with pullorum or typhoid are quarantined until marketed or destroyed under the supervision of the Official State Agency, or until subsequently blood tested following the procedure for reacting flocks as contained in §145.14(a)(5), and all birds fail to demonstrate pullorum or typhoid infection;

(b)(3)(vii) All poultry, including exhibition, exotic, and game birds, but excluding waterfowl, going to public exhibition shall come from U.S. Pullorum-Typhoid Clean or equivalent flocks, or have had a negative pullorum-typhoid test within 90 days of going to public exhibition;

(b)(3)(viii) Discontinuation of any of the conditions or procedures described in paragraphs (b)(3)(i), (ii), (iii), (iv), (v), (vi), and (vii) of this section, or the occurrence of repeated outbreaks of pullorum or typhoid in poultry breeding flocks, other than turkey flocks, within or originating within the State shall be grounds for the Service to revoke its determination that such conditions and procedures have been met or complied with. Such action shall not be taken until a thorough investigation has been made by the Service and the Official State Agency has been given an opportunity to present its views.

(b)(4) It is a multiplier breeding flock located in a State which has been determined by the Service to be in compliance with the provisions of paragraph (b)(3) of this section, and in which pullorum disease or fowl typhoid is not known to exist nor to have existed in hatchery supply flocks, other than turkey, waterfowl, exhibition poultry, and game bird supply flocks, within the State during the preceding 12 months.

(b)(5) It is a primary breeding flock located in a State determined to be in compliance with the provisions of paragraph (b)(4) of this section, and in which a sample of 300 birds from flocks of more than 300, and each bird in flocks of 300 or less, has been officially tested for pullorum-typhoid with no reactors: Provided, That a bacteriological examination monitoring program acceptable to the Official State Agency and approved by the Service may be used in lieu of blood testing.

(c) U.S. M. Gallisepticum Clean. (1) A flock maintained in compliance with the provisions of §147.26 of this chapter and in which freedom from M. gallisepticum has been demonstrated under the criteria specified in paragraph (c)(1)(i) or (ii) of this section.

(c)(1)(i) It is flock in which all birds or a sample of at least 300 birds has been tested for M. gallisepticum as provided in §145.14(b) when more than 4 months of age: Provided, That to retain this classification, a minimum of 150 birds shall be tested at intervals of not more than 90 days: And provided further, That a sample comprised of less than 150 birds may be tested at any one time, with the approval of the Official State Agency and the concurrence of the Service, provided that a minimum of 150 birds is tested within each 90-day period; or

(c)(1)(ii) It is a multiplier breeding flock which originated as U.S. M. Gallisepticum Clean chicks from primary breeding flocks and from which a sample comprised of a minimum of 150 birds per flock has been tested for M. gallisepticum as provided in §145.14(b) when more than 4 months of age: Provided, That to retain this classification, the flock shall be subjected to one of the following procedures:

(c)(1)(a) At intervals of not more than 90 days, a sample 75 birds, with a minimum of 30 birds per pen, whichever is greater, shall be tested; or

(c)(1)(b) At intervals of not more than 30 days, a sample of 25 cull chicks produced from the flock shall be subjected to laboratory procedures acceptable to the Official State Agency and approved by the Service, for the detection and recovery of M. gallisepticum.

(c)(2) A participant handling U.S. M. Gallisepticum Clean products shall keep these products separate from other products in a manner satisfactory to the Official State Agency: Provided, That U.S. M. Gallisepticum Clean chicks from primary breeding flocks shall be produced in incubators and hatchers in which only eggs from flocks qualified under paragraph (c)(1)(i) of this section are set.

(c)(3) U.S. M. Gallisepticum Clean chicks shall be boxed in clean boxes and delivered in trucks that have been cleaned and disinfected as described in §147.24(a) of this chapter.

(d) U.S. Sanitation Monitored. This program is intended to be the basis from which the breeding-hatching industry may conduct a program for the prevention and control of Salmonellosis. It is intended to reduce the incidence of Salmonella organisms in hatching eggs and chicks through an effective and practical sanitation program at the breeder farm and in the hatchery. This will afford other segments of the poultry industry an opportunity to reduce the incidence of Salmonella in their products.

(d)(1) A flock and the hatching eggs and chicks produced from it which have met the following requirements as determined by the Official State Agency:

(d)(1)(i) The flock shall originate from a source where sanitation and management practices, as outlined in §145.33(d)(1) of this paragraph, are conducted;

(d)(1)(ii) The flock is maintained in compliance with §§147.21, 147.24(a), and 147.26 of this chapter;

(d)(1)(iii) All feed should be pelletized and/or crumbled in mills operated at 190°F. or above in facilities for recontamination control;

(d)(1)(iv) Feed shall be stored and transported in such a manner as to prevent possible contamination;

(d)(1)(v) Hatching eggs are collected at least four times a day and are handled as described in §147.22 and fumigated on the farm as described in §147.25(a) of this chapter or handled in a sanitary manner equivalent to §§147.22 and 147.25(a) of this part, which has been approved by the Official State Agency with the concurrence of the Service; and

(d)(1)(vi) Chicks are hatched in a hatchery meeting the requirements of §§147.23 and 147.24(b) and are in compliance with §147.25(b) of this chapter.

(d)(2) The Official State Agency may use the procedures described in §147.14 of this chapter to monitor the effectiveness of the sanitation practices.

(d)(3) In order for a hatchery to sell products of this classification, all products handled shall meet the requirements of the classification.

(d)(4) This classification may be revoked by the Official State Agency if the participant fails to follow recommended corrective measures.

(e) U.S. M. Synoviae Clean. (1) A flock maintained in compliance with the provisions of §147.26 and in which freedom from M. synoviae has been demonstrated under the criteria specified in paragraph (e)(1)(i) or (ii) of this section.

(e)(1)(i) It is a flock in which a minimum of 300 birds has been tested for M. synoviae as provided for in §145.14(b) when more than 4 months of age: Provided, That to retain this classification, a sample of at least 150 birds shall be tested at intervals of not more than 90 days: And provided further, That a sample comprised of less than 150 birds may be tested at any one time, with the approval of the Official State Agency and the concurrence of the Service, provided that a minimum of 150 birds is tested within each 90-day period; or

(e)(1)(ii) It is a multiplier breeding flock which originated as U.S. M. Synoviae Clean chicks from primary breeding flocks and from which a sample comprised of a minimum of 75 birds has been tested for M. synoviae as provided in §145.14(b) when more than 4 months of age: Provided, That to retain this classification, a sample of 50 birds shall be tested at intervals of not more than 90 days: And provided further, That a sample of less than 50 birds may be tested at any one time, provided that a minimum of 30 birds per flock with a minimum of 15 birds per pen, whichever is greater, is tested each time and a total of at least 50 birds is tested within each 90-day period.

(e)(2) A participant handling U.S. M. Synoviae Clean products shall keep these products separate from other products in a manner satisfactory to the Official State Agency: Provided, That U.S. M. Synoviae Clean chicks from primary breeding flocks shall be produced in incubators and hatchers in which only eggs from flocks qualified under paragraph (e)(1)(i) or (ii) of this section are set.

(e)(3) U.S. M. Synoviae Clean chicks shall be boxed in clean boxes and delivered in trucks that have been cleaned and disinfected as described in §147.24(a).

§145.34 Terminology and classification; States.

(a) U.S. Pullorum-Typhoid Clean State. (1) A State will be declared a U.S. Pullorum-Typhoid Clean State when it has been determined by the Service that:

(a)(1)(i) The State is in compliance with the provisions contained in §145.23(b)(3)(i) through (vii), §145.33(b)(3)(i) through (vii), §145.43(b)(3)(i) through (vi), and §145.53(b)(3)(i) through (vii).

(a)(1)(ii) No pullorum disease or fowl typhoid is known to exist nor to have existed in hatchery supply flocks within the State during the preceding 12 months: Provided, That pullorum disease or fowl typhoid found in waterfowl, exhibition poultry, and game bird breeding flocks will not prevent a State, which is otherwise eligible, from qualifying for a period of 2 years.

(a)(2) Discontinuation of any of the conditions described in paragraph (a)(1)(i) of this section, or repeated outbreaks of pullorum or typhoid occur in hatchery supply flocks described in paragraph (a)(1)(ii) of this section, or if an infection spreads from the originating premises, the Service shall have grounds to revoke its determination that the State is entitled to this classification. Such action shall not be taken until a thorough investigation has been made by the Service and the Official State Agency has been given an opportunity for a hearing.

Subpart D - Special Provisions For Turkey Breeding Flocks And Products

§145.41 Definitions.

Except where the context otherwise requires, for the purposes of this subpart the following terms shall be construed respectively, to mean:

(a) Poults. Newly hatched turkeys which have not been fed or watered.

§145.42 Participation.

(a) Participating turkey flocks, and the eggs and poults produced from them, shall comply with the applicable general provisions of Subpart A of this part and the special provisions of this Subpart D.

(b) The minimum weight of turkey hatching eggs shipped interstate shall be 2 ounces each for small varieties and 2 1/2 ounces each for other varieties, unless otherwise specified by the purchaser of the eggs.

(c) Hatching eggs shall be fumigated according to the procedures described in §147.25(a): Provided, That alternative sanitizing procedures may be used with the approval of the Official State Agency in each specific instance and with the general concurrence by the Service in the policy adopted by the Official State Agency.

§145.43 Terminology and classification; flocks and products.

Participating flocks, and the eggs and poults produced from them, which have met the respective requirements specified in this section may be designated by the following terms and the corresponding designs illustrated in §145.10:

(a) [Reserved]

(b) U.S. Pullorum-Typhoid Clean. A flock in which freedom from pullorum and typhoid has been demonstrated to the Official State Agency under the criteria in one of the following paragraphs (b)(1) through (5) of this section: Provided, That a flock qualifying by means of a blood test shall be tested within the past 12 months, except that the retesting of a participating flock which is retained for more than 12 months shall be at the discretion of the Official State Agency with the concurrence of the Service. (See §145.14 relating to the official blood test where applicable.)

(b)(1) It has been officially blood tested with no reactors.

(b)(2) It is a multiplier breeding flock, or a breeding flock composed of progeny of a primary breeding flock which is intended solely for the production of multiplier breeding flocks, and meets the following specifications as determined by the Official State Agency and the Service:

(b)(2)(i) The flock is located in a State where all persons performing poultry disease diagnostic services within the State are required to report to the Official State Agency within 48 hours the source of all poultry specimens from which S. pullorum or S. gallinarum is isolated;

(b)(2)(ii) The flock is composed entirely of birds that originated from U.S. Pullorum-Typhoid Clean breeding flocks or from flocks that met equivalent requirements under official supervision; and

(b)(2)(iii) A sample comprised of at least 25 percent of the birds in the flock has been officially blood tested with no reactors, or its progeny has been subjected to an approved 10-day poult mortality bacteriological examination monitoring program and bacteriological examination of a sample of down shed by poults in the hatcher from selected hatches as prescribed by the Official State Agency: Provided, That when the blood testing procedure is used the percentage of the flock included in the sample may be reduced by 5 percentage points following each year in which there is no evidence of infection on the premises until the required percentage is reduced to zero: And provided further, That the sample tested for the qualification of a flock under this subparagraph shall include at least 500 birds the first year, 400 the second year, 300 the third year, 200 the fourth year, and 100 the fifth year.

(b)(3) It is a multiplier breeding flock, or a breeding flock composed of progeny of a primary breeding flock which is intended solely for the production of multiplier breeding flocks, that originated from U.S. Pullorum-Typhoid Clean breeding flocks or from flocks that met equivalent requirements under official supervision, and is located in a State in which it has been determined by the Service that:

(b)(3)(i) All turkey hatcheries within the State are qualified as "National Plan Hatcheries" or have met equivalent requirements for pullorum-typhoid control under official supervision;

(b)(3)(ii) All turkey hatchery supply flocks within the State are qualified as U.S. Pullorum-Typhoid Clean or have met equivalent requirements for pullorum-typhoid control under official supervision: Provided, That if other domesticated fowl, except waterfowl, are maintained on the same premises as the participating flock, freedom from pullorum-typhoid infection shall be demonstrated by an official blood test of each of these fowl;

(b)(3)(iii) All shipments of products other than U.S. Pullorum-Typhoid Clean, or equivalent, into the State are prohibited;

(b)(3)(iv) All persons performing poultry disease diagnostic services within the State are required to report to the Official State Agency within 48 hours the source of all poultry specimens from which S. pullorum or S. gallinarum is isolated;

(b)(3)(v) All reports of S. pullorum or S. gallinarum isolations from poultry are promptly followed by an investigation by the Official State Agency to determine the origin of the infection;

(b)(3)(vi) All flocks found to be infected with pullorum or typhoid are quarantined until marketed or destroyed under the supervision of the Official State Agency, or until subsequently blood tested, following the procedure for reacting flocks as contained in §145.14(a)(5), and all birds fail to demonstrate pullorum or typhoid infection;

(b)(3)(vii) [Reserved]

(b)(3)(viii) Discontinuation of any of the conditions or procedures described in paragraphs (b)(3)(i), (ii), (iii), (iv), (v), and (vi) of this section, or the occurrence of repeated outbreaks of pullorum or typhoid in turkey breeding flocks within or originating within the State shall be grounds for the Service to revoke its determination that such conditions and procedures have been met or complied with. Such action shall not be taken until a thorough investigation has been made by the Service and the Official State Agency has been given an opportunity to present its views.

(b)(4) It is a multiplier breeding flock located in a State which has been determined by the Service to be in compliance with the provisions of paragraph (b)(3) of this section and in which pullorum disease or fowl typhoid is not known to exist nor to have existed in turkey hatchery supply flocks within the State during the preceding 24 months.

(b)(5) It is a primary breeding flock located in a State determined to be in compliance with the provisions of paragraph (b)(4), of this section and in which a sample of 300 birds from flocks of more than 300, and each bird in flocks of 300 or less, has been officially tested for pullorum-typhoid with no reactors: Provided, That a bacteriological examination monitoring program acceptable to the Official State Agency and approved by the Service may be used in lieu of blood testing.

(c) U.S. M. Gallisepticum Clean. (1) A flock maintained in accordance with the conditions and procedures described in §147.26 of this chapter, and in which no reactors are found when a random sample of at least 10 percent of the birds in the flock, or 300 birds in flocks of more than 300 and each bird in flocks of 300 or less, is tested when more than 4 months of age, in accordance with the procedures described in §145.14(b).

(c)(2) A flock qualified as U.S. M. Gallisepticum Clean may retain the classification for 1 year, provided it is maintained in isolation and no evidence of M. gallisepticum infection is revealed. Each flock and premises shall be inspected at least once during the laying period by an Authorized Agent of the Official State Agency or the State Animal Disease Control Official. If a flock proves to be infected with M. gallisepticum, it shall lose this classification.

(c)(3) In order to sell hatching eggs or poults of this classification, all hatching eggs and poults handled by the participant must be of this classification.

(d) U.S. M. Meleagridis Clean.[1/] (1) A flock in which freedom from M. meleagridis has been demonstrated under the following criteria:

(d)(1)(i) A sample of 60 birds from each flock has been tested for M. meleagridis when more than 4 months of age: Provided, That to retain this classification, a minimum of 30 samples from male flocks and 60 samples from female flocks shall be retested at 28-30 weeks of age and at 4-6 week intervals thereafter.

(d)(2) The official blood tests for M. meleagridis shall be the serum plate agglutination, the tube agglutination, or the microagglutination tests. The hemagglutination inhibition tests, serum plate dilution test, and microagglutination tests may be used as supplemental tests to determine the status of the flock, in accordance with §147.6(b).

(d)(3) The tests shall be conducted using M. meleagridis antigens and the protocols for testing approved by the Department or the Official State Agency.

1/ Effective Date: January 1, 1983.

(d)(4) When reactors to the official test are found and can be identified, 10 tracheal swabs and/or vaginal or phallus swabs and their corresponding blood samples shall be submitted to a laboratory for serological and cultural examination. If reactors cannot be identified, at least 30 tracheal swabs and/or vaginal or phallus swabs and their corresponding blood samples shall be submitted. In a flock with a low reactor rate (less than 5 reactors) the reactors may be submitted to the laboratory within 10 days for serology, necropsy, and thorough bacteriological examination.

(d)(5) If a mycoplasma is isolated, the organism must be serotyped. If M. meleagridis is isolated, the flock shall be considered infected.

§145.44 Terminology and classification; States.

(a) U.S. Pullorum-Typhoid Clean State. (1) A State will be declared a U.S. Pullorum-Typhoid Clean State when it has been determined by the Service that:

(a)(1)(i) The State is in compliance with the provisions contained in §145.23(b)(3)(i) through (vii), §145.33(b)(3)(i) through (vii), §145.43(b)(3)(i) through (vi), and §145.53(b)(3)(i) through (vii).

(a)(1)(ii) No pullorum disease or fowl typhoid is known to exist nor to have existed in hatchery supply flocks within the State during the preceding 12 months: Provided, That pullorum disease or fowl typhoid found in waterfowl, exhibition poultry, and game bird breeding flocks will not prevent a State, which is otherwise eligible, from qualifying for a period of 2 years.

(a)(2) Discontinuation of any of the conditions described in paragraph (a)(1)(i) of this section, or repeated outbreaks of pullorum or typhoid occur in hatchery supply flocks described in paragraph (a)(1)(ii) of this section, or if an infection spreads from the originating premises, the Service shall have grounds to revoke its determination that the State is entitled to this classification. Such action shall not be taken until a thorough investigation has been made by the Service and the Official State Agency has been given an opportunity for a hearing.

(b) U.S. Pullorum-Typhoid Clean State, Turkeys. (1) A State will be declared a U.S Pullorum-Typhoid Clean State, Turkeys, when it has been determined by the Service that:

(b)(1)(i) The State is in compliance with the provisions contained in §145.43(b)(3)(i) through (vi).

(b)(1)(ii) No pullorum disease or fowl typhoid is known to exist nor to have existed in turkey hatchery supply flocks within the State during the preceding 24 months.

(b)(2) Discontinuation of any of the conditions described in paragraph (b)(1)(i) of this section, or repeated outbreaks of pullorum or typhoid occur in hatchery supply flocks described in paragraph (b)(1)(ii) of this section, or if an infection spreads from the originating premises, the Service shall have grounds to revoke its determination that the State is entitled to this classification. Such action shall not be taken until a thorough investigation has been made by the Service and the Official State Agency has been given an opportunity for a hearing.

(c) U.S. M. Gallisepticum Clean State, Turkeys. (1) A State will be declared a U.S. M. Gallisepticum Clean State, Turkeys, when it has been determined by the Service that:

(c)(1)(i) No M. gallisepticum is known to exist nor to have existed in turkey breeding flocks in production within the State during the preceding 12 months;

(c)(1)(ii) All turkey breeding flocks in production are classified as U.S. M. Gallisepticum Clean or have met equivalent requirements for M. gallisepticum control under official supervision;

(c)(1)(iii) All turkey hatcheries within the State handle products which are classified as U.S. M. Gallisepticum Clean or have met equivalent requirements for M. gallisepticum control under official supervision;

(c)(1)(iv) All shipments of turkey products other than those classified as U.S. M. Gallisepticum Clean, or equivalent, into the State are prohibited;

(c)(1)(v) All persons performing poultry disease diagnostic services within the State are required to report to the Official State Agency within 48 hours the source of all turkey specimens that have been identified as being infected with M. gallisepticum;

(c)(1)(vi) All reports of M. gallisepticum infection in turkeys are promptly followed by an investigation by the Official State Agency to determine the origin of the infection;

(c)(1)(vii) All turkey flocks found to be infected with M. gallisepticum are quarantined until marketed under supervision of the Official State Agency.

(c)(2) Discontinuation of any of the conditions described in paragraph (c)(1) of this section, or if repeated outbreaks of M. gallisepticum occur in turkey breeding flocks described in paragraph (c)(1)(ii) of this section, or if an infection spreads from the originating premises, the Service shall have grounds to revoke its determination that the State is entitled to this classification. Such action shall not be taken until a thorough investigation has been made by the Service and the Official State Agency has been given an opportunity for a hearing.

Subpart E - Special Provisions For Waterfowl, Exhibition Poultry, And Game Bird Breeding Flocks And Products

§145.51 Definitions.

Except where the context otherwise requires, for the purposes of this subpart the following terms shall be construed, respectively, to mean:

(a) Waterfowl. Domesticated fowl that normally swim, such as ducks and geese.

(b) Exhibition Poultry. Domesticated fowl which are bred for the combined purposes of meat or egg production and competitive showing.

(c) Game birds. Domesticated fowl such as pheasants, partridge, quail, grouse, and guineas, but not doves and pigeons.

§145.52 Participation.

Participating flocks of waterfowl, exhibition poultry, and game birds, and the eggs and baby poultry produced from them shall comply with the applicable general provisions of Subpart A of this part and the special provisions of this Subpart E.

(a) Started poultry shall lose their identity under Plan terminology when not maintained by Plan participants under the conditions prescribed in §145.5(a).

(b) Hatching eggs produced by primary breeding flocks shall be fumigated according to the procedures described in §147.25(a) of this chapter: Provided, That alternative sanitizing procedures may be used with the approval of the Official State Agency in each specific instance and with the general concurrence by the Service in the policy adopted by the Official State Agency.

§145.53 Terminology and classification; flocks and products.

Participating flocks, and the eggs and baby poultry produced from them, which have met the respective requirements specified in this section may be designated by the following terms and the corresponding designs illustrated in §145.10.

(a) U.S. Approved. All birds in the breeding flock observed by Authorized Agents or State Inspectors are found to conform with the criteria for the breed represented, as contained in the Standard of Perfection[1/] or the breeder's specifications for the stock represented in the flock, and such specifications are on file with the Official State Agency.

(b) U.S. Pullorum-Typhoid Clean. A flock in which freedom from pullorum and typhoid has been demonstrated to the Official State Agency under the criteria in one of the following paragraphs (b)(1) through (5) of this section. (See §145.14 relating to the official blood test where applicable.):

(b)(1) It has been officially blood tested within the past 12 months with no reactors.

(b)(2) It is a multiplier breeding flock, or a breeding flock composed of progeny of a primary breeding flock which is intended solely for the production of multiplier breeding flocks, and meets the following specifications as determined by the Official State Agency and the Service:

(b)(2)(i) The flock is located in a State where all persons performing poultry disease diagnostic services within the State are required to report to the Official State Agency within 48 hours the source of all poultry specimens from which S. pullorum or S. gallinarum is isolated;

(b)(2)(ii) The flock is composed entirely of birds that originated from U.S. Pullorum-Typhoid Clean breeding flocks or from flocks that met equivalent requirements under official supervision; and

(b)(2)(iii) A sample comprised of at least 25 percent of the birds in the flock has been officially blood tested within the past 12 months with no reactors, or its progeny has been subjected to an approved 10-day baby poultry mortality bacteriological examination monitoring program and bacteriological examination of a sample of down shed by baby poultry in the hatcher from selected hatches as prescribed by the Official State Agency: Provided, That when the blood testing procedure is used, the percentage of the flock included in the sample may be reduced by 5 percentage points following each year in which there is no evidence of infection on the premises until the required percentage is reduced to zero: And provided further, That the sample tested for the qualification of a flock under this subparagraph shall include at least 500 birds the first year, 400 the second year, 300 the third year, 200 the fourth year, and 100 the fifth year.

(b)(3) It is a multiplier breeding flock, or a breeding flock composed of progeny of a primary breeding flock which is intended solely for the production of multiplier breeding flocks, that originated from U.S. Pullorum-Typhoid Clean breeding flocks or from flocks that met equivalent requirements under official supervision, and is located in a State in which it has been determined by the Service that:

[1/] Published by the American Poultry Association, Inc.

(b)(3)(i) All hatcheries, except turkey hatcheries, within the State are qualified as "National Plan Hatcheries" or have met equivalent requirements for pullorum-typhoid control under official supervision;

(b)(3)(ii) All hatchery supply flocks, except turkey flocks, within the State, are qualified as U.S. Pullorum-Typhoid Clean or have met equivalent requirements for pullorum-typhoid control under official supervision: Provided; That if other domesticated fowl, except waterfowl, are maintained on the same premises as the participating flock, freedom from pullorum-typhoid infection shall be demonstrated by an official blood test of each of these fowl;

(b)(3)(iii) All shipments of products other than U.S. Pullorum-Typhoid Clean, or equivalent, into the State are prohibited;

(b)(3)(iv) All persons performing poultry disease diagnostic services within the State are required to report to the Official State Agency within 48 hours the source of all poultry specimens from which S. pullorum or S. gallinarum is isolated;

(b)(3)(v) All reports of S. pullorum or S. gallinarum isolations from poultry are promptly followed by an investigation by the Official State Agency to determine the origin of the infection;

(b)(3)(vi) All flocks found to be infected with pullorum or typhoid are quarantined until marketed or destroyed under the supervision of the Official State Agency, or until subsequently blood tested, following the procedure for reacting flocks as contained in §145.14(a)(5), and all birds fail to demonstrate pullorum or typhoid infection;

(b)(3)(vii) All poultry, including exhibition, exotic, and game birds, but excluding waterfowl, going to public exhibition shall come from U.S. Pullorum-Typhoid Clean or equivalent flocks, or have had a negative pullorum-typhoid test within 90 days of going to public exhibition;

(b)(3)(viii) Discontinuation of any of the conditions or procedures described in paragraphs (b)(3)(i), (ii), (iii), (iv), (v), (vi), and (vii) of this section, or the occurrence of repeated outbreaks of pullorum or typhoid in poultry breeding flocks, other than turkey flocks, within or originating within the State shall be grounds for the Service to revoke its determination that such conditions and procedures have been met or complied with. Such action shall not be taken until a thorough investigation has been made by the Service and the Official State Agency has been given an opportunity to present its views.

(b)(4) It is a multiplier breeding flock located in a State which has been determined by the Service to be in compliance with the provisions of paragraph (b)(3) of this section, and in which pullorum disease or fowl typhoid is not known to exist nor to have existed in hatchery supply flocks, other than turkey flocks, within the State during the preceding 24 months.

(b)(5) It is a primary breeding flock located in a State determined to be in compliance with the provisions of paragraph (b)(4) of this section, and in which a sample of 300 birds from flocks of more than 300, and each bird in flocks of 300 or less, has been officially tested for pullorum-typhoid within the past 12 months with no reactors: Provided, That a bacteriological examination monitoring program or serological examination monitoring program for game birds acceptable to the Official State Agency and approved by the Service may be used in lieu of annual blood testing.

(c) U.S. M. Gallisepticum Clean. (1) A flock maintained in compliance with the provisions of §147.26 of this chapter and in which freedom from M. gallisepticum has been demonstrated under the criteria specified in paragraph (c)(1)(i) or (ii) of this section.

(c)(1)(i) It is a flock in which all birds or a sample of at least 300 birds has been tested for M. gallisepticum as provided in §145.14(b) when more than 4 months of age or upon reaching sexual maturity: Provided, That to retain this classification, a sample of at least 5 percent of the birds in the flock, with a minimum of 100 birds, shall be tested at intervals of not more than 90 days: And provided further, That a sample comprised of less than 5 percent may be tested at any one time, with the approval of the Official State Agency and the concurrence of the Service, provided that a total of at least 5 percent of the birds in the flock, with a minimum of 100 birds, is tested within each 90-day period; or

(c)(1)(ii) It is a multiplier breeding flock which originated as U.S. M. Gallisepticum Clean baby poultry from primary breeding flocks and a random sample comprised of 50 percent of the birds in the flock, with a maximum of 200 birds and a minimum of 30 birds per flock, has been tested for M. gallisepticum as provided in §145.14(b) when more than 4 months of age or upon reaching sexual maturity: Provided, That to retain this classification, the flock shall be subjected to one of the following procedures:

(c)(1)(ii)(a) At intervals of not more than 90 days, a random sample of at least 2 percent of the birds in the flock, with a minimum of 30 birds per pen, shall be tested; or

(c)(1)(ii)(b) At intervals of not more than 30 days, a sample of 25 cull baby poultry produced from the flock shall be subjected to laboratory procedures acceptable to the Official State Agency and approved by the Service, for the detection and recovery of M. gallisepticum.

(c)(2) A participant handling U.S. M. Gallisepticum Clean products shall keep these products separate from other products in a manner satisfactory to the Official State Agency: Provided, That U.S. M. Gallisepticum Clean baby poultry from primary breeding flocks shall be produced in incubators and hatchers in which only eggs from flocks qualified under paragraph (c)(1)(i) of this section are set.

(c)(3) U.S. M. Gallisepticum Clean baby poultry shall be boxed in clean boxes and delivered in trucks that have been cleaned and disinfected as described in §147.24(a) of this chapter.

§145.54 Terminology and classification; States.

(a) U.S. Pullorum-Typhoid Clean State. (1) A State will be declared a U.S. Pullorum-Typhoid Clean State when it has been determined by the Service that:

(a)(1)(i) The State is in compliance with the provisions contained in §145.23(b)(3)(i) through (vii), §145.33(b)(3)(i) through (vii), §145.43(b)(3)(i) through (vi), and §145.53(b)(3)(i) through (vii).

(a)(1)(ii) No pullorum disease or fowl typhoid is known to exist nor to have existed in hatchery supply flocks within the State during the preceding 12 months: Provided, That pullorum disease or fowl typhoid found in waterfowl, exhibition poultry, and game bird breeding flocks will not prevent a State, which is otherwise eligible, from qualifying for a period of 2 years.

(a)(2) Discontinuation of any of the conditions described in paragraph (a)(1)(i) of this section, or repeated outbreaks of pullorum or typhoid occur in hatchery supply flocks described in paragraph (a)(1)(ii) of this section, or if an infection spreads from the originating premises, the Service shall have grounds to revoke its determination that the State is entitled to this classification. Such action shall not be taken until a thorough investigation has been made by the Service and the Official State Agency has been given an opportunity for a hearing.

PART 146 - [Reserved]

PART 147 - AUXILIARY PROVISIONS ON NATIONAL POULTRY IMPROVEMENT PLAN

Subpart A - Blood Testing Procedures

Sec.
147.1 The standard tube agglutination test.
147.2 The rapid serum test.
147.3 The stained-antigen, rapid, whole-blood test.
147.4 The tube agglutination test for S. typhimurium.
147.5 The microagglutination test for pullorum-typhoid.
147.6 Procedure for determining the status of flocks reacting to test for Mycoplasma gallisepticum and Mycoplasma synoviae.

Subpart B - Bacteriological Examination Procedures

147.11 Laboratory procedure recommended for the bacteriological examination of salmonella reactors.
147.12 Procedures for collecting environmental samples and cloacal swabs for bacteriological examination.
147.13 Procedure for bacteriological culturing of eggshells for colon bacilli organisms.
147.14 Procedures to determine status and effectiveness of sanitation monitored program.
147.15 Laboratory procedure recommended for the bacteriological examination of mycoplasma reactors.
Sec.
147.16 Procedure for the evaluation of mycoplasma reactors by in vivo bioassay (enrichment).

Subpart C - Sanitation Procedures

147.21 Flock sanitation.
147.22 Hatching egg sanitation.
147.23 Hatchery sanitation.
147.24 Cleaning and disinfecting.
147.25 Fumigation.
147.26 Procedures for establishing isolation and maintaining sanitation and good management practices for the control of Salmonella and Mycoplasma infections.
147.27 Procedures recommended to prevent the spread of disease by artificial insemination of turkeys.

Subpart D - [Reserved]

Subpart E - Procedure for Changing National Poultry Improvement Plan

147.41 Definitions.
147.42 General.
147.43 General Conference Committee.
147.44 Submitting, compiling, and distributing proposed changes.
147.45 Official delegates.
147.46 Committee consideration of proposed changes.

Sec.
147.47 Conference consideration of proposed changes.
147.48 Approval of conference recommendations by the Department.

AUTHORITY: Sec. 101(b) of the Department of Agriculture Organic Act of 1944, as amended, 58 Stat. 734 (7 U.S.C. 429).

Subpart A - Blood Testing Procedures

§147.1 The standard tube agglutination test.[1/]

(a) The blood samples should be collected and delivered as follows:

(a)(1) The blood samples should be taken by properly qualified and authorized persons only, and in containers provided by the laboratory. The containers should be stout-walled test tubes, preferably 3/8 by 3 inches, without lip, or small well-selected medicine vials, which have been thoroughly cleaned and dried in a hot-air drying oven. If stoppers are used, they should be thoroughly cleaned and dried.

(a)(2) Sufficient blood should be procured by making a small incision in the large median wing vein with a small sharp lancet and allowing the blood to run into the tube, or by the use of a small syringe (with 20 or 21 gage needle) which is properly cleansed between bleedings with physiological saline solution. To facilitate the separation of the serum, the tubes should be placed in a slanted position until the blood has solidified. After the blood has completely clotted, they should be packed and shipped by mail (special delivery), rapid express, or by messenger, to the laboratory. All labeling must be clear and permanent, and may be done with a suitable pencil on etched portions of the tube, or by means of fast-gum labels.

(a)(3) The blood samples must reach the laboratory in a fresh and unhemolyzed condition. Hemolyzed samples should be rejected. It is imperative, therefore, to cool the tubes immediately after slanting and clotting, and unless they reach the laboratory within a few hours, to pack them with ice in special containers, or use some other cooling system which will insure their preservation during transportation. In severe cold seasons, extreme precautions must be exercised to prevent freezing and consequent laking. The samples must be placed in cold (5° to 10°C) storage, immediately upon arrival at the laboratory.

(b) The antigen shall consist of representative strains of S. pullorum which are of known antigenic composition, high agglutinability, but are not sensitive to negative and nonspecific sera. The stock cultures may be maintained satisfactorily by transferring to new sloped agar at least once a month and keeping at 18° to 25°C (average room temperature) in a dark closet or chest, following incubation for from 24 to 36 hours at 37°C. The antigenic composition and purity of the stock cultures should be checked consistently.

[1/] The procedure described is a modification of the method reported in the proceedings of the U.S. Livestock Sanitary Association, November 30 to December 2, 1932, pp-487 to 491.

(c) A medium which has been used satisfactorily has the following composition:

Water	1,000 cc
Difco beef extract	4 gm. (0.4 percent)
Difco Bacto-peptone	10 gm. (1.0 percent)
Difco dry-granular agar	20 gm. (2.0 percent)

Reaction-pH 6.8 to 7.2.

(d) Large 1-inch test tubes, Kolle flasks, or Blake bottles should be streaked liberally over the entire agar surface with inoculum from 48-hour slant agar cultures prepared from the stock cultures of the selected strains. The antigen-growing tubes or bottles should be incubated 48 hours at 37°C, and the surface growth washed off with sufficient phenolized (0.5 percent) saline (0.85 percent) solution to make a heavy suspension. The suspension should be filtered free of clumps through a thin layer of absorbent cotton in a Buchner funnel with the aid of suction. The antigens of the separate strains should be combined in equal volume-density and stored in the refrigerator (5° to 10°C) in tightly stoppered bottles.

(e) Thiosulfate-Glycerin (TG) medium may be used as an alternate medium for the preparation of tube agglutination antigen. The TG medium, formerly used for the preparation of stained, whole-blood antigen, is described in more detail in the article by A.D. MacDonald, Recent Developments in Pullorum Antigen for the Rapid, Whole-Blood Test, Report of the Conference of the National Poultry Improvement Plan, pages 122-127, 1941. This medium provides a tube antigen of excellent specificity and greatly increases the yield of antigen from a given amount of medium. The TG medium has the following composition:

Beef infusion	1,000 cc
Difco Bacto-peptone	20 gm. (2.0 percent)
Sodium Thiosulfate	5 gm. (0.5 percent)
Ammonium chloride	5 gm. (0.5 percent)
Glycerin, U.S.P. (95 percent)	20 cc (2.0 percent)
Difco dry-granular agar	30 gm. (3.0 percent)

Reaction-pH 6.8 to 7.2.

Large 1-inch test tubes, Kolle flasks, Blake bottles, or Erlenmeyer flasks should be seeded over the entire agar surface with inoculum from 24-hour beef infusion broth cultures prepared from the stock cultures of the selected strains. The antigen-growing tubes or bottles should be incubated 96 hours at 37°C, and the surface growth washed off with sufficient phenolized (0.5 percent) saline (0.85 percent) solution to make a heavy suspension. The suspension should be filtered free of clumps through a thin layer of absorbent cotton in a Buchner funnel with the aid of suction. The antigen then should be centrifuged. The mass of bacteria should be removed from the centrifuge tubes or bowl and resuspended in saline (0.85 percent) solution containing 0.5 percent phenol. After the bacterial mass has been uniformly suspended in the diluent, it should be again passed through a cotton pad in a Buchner funnel without the aid of suction. The antigens of the separate strains should be combined in equal volume-density and stored in the refrigerator (5° to 10°C) in tightly stoppered bottles.

(f) The diluted antigen to be used in the routine testing should be prepared from the stock antigen by dilution of the latter with physiological (0.85 percent) saline solution containing 0.25 percent of phenol to a turbidity corresponding to 0.75-1.00 on the McFarland nephelometer scale. The hydrogen-ion concentration of the diluted antigen should be corrected to pH 8.2 to 8.5 by the addition of dilute sodium hydroxide. New diluted antigen should be prepared each day and kept cold. The diluted antigen may be employed in 2 cc quantities in 4 by 1/2-inch test tubes, or 1 cc quantities in smaller tubes, in which the final serum-antigen mixtures are made and incubated. The distribution of the antigen in the tubes may be accomplished by the use of long burettes, or special filling devices made for the purpose.

(g) The maximum serum dilution employed must not exceed 1:50 for chickens, nor 1:25 for turkeys. The available data indicate that 1:25 dilution is the most efficient. In all official reports on the blood test, the serum dilutions shall be indicated. The sera should be introduced into the agglutination tubes in the desired amounts with well-cleaned serological pipettes or special serum-delivery devices which do not permit the mixing of different sera. The antigen and serum should be well mixed before incubation. The serum and antigen mixture must be incubated for at least 20 hours at 37°C.

(h) The results shall be recorded as:

N, or - (negative) when the serum-antigen mixture remains uniformly turbid.

P, or + (positive) when there is a distinct clumping of the antigen, and the liquid between the agglutinated particles is clear.

S, or ? (suspicious) when the agglutination is only partial or incomplete.

M, or missing, when samples listed on the original record sheet are missing.

H, or hemolyzed, when blood samples are hemolyzed and connot be tested.

B, or broken, when sample tubes are broken and no serum can be obtained.

(Some allowance must always be made for the difference in sensitiveness of different antigens and different set-ups, and therefore, a certain amount of independent, intelligent judgment must be exercised at all times. Also, the histories of the flocks require consideration. In flocks where individuals show a suspicious agglutination, it is desirable to examine representative birds bacteriologically to determine the presence or absence of S. pullorum.)

§147.2 The rapid serum test.[2/]

(a) The procedure for the collection and delivery of blood samples in the rapid serum test is the same as that described in §147.1(a).

(b) The selection and maintenance of suitable strains of S. pullorum and the composition of a satisfactory medium are described in §147.1 (b) and (c).

(c) Large 1-inch test tubes, Kolle flasks, or Blake bottles are streaked liberally from 48-hour slant-agar cultures prepared from stock cultures of the selected strains.

2/ The procedure described is a modification of the method reported by Runnells, Coon, Farley, and Thorpe, Amer. Vet. Med. Assoc. Jour. 70 (N.S. 23):660-662 (1927).

(d) The antigen-growing tubes or bottles should be incubated 48 hours at 37°C, and the surface growth washed off with a very slight amount of 12 percent solution of sodium chloride containing 0.25 to 0.5 percent phenol, filtered through lightly packed sterile absorbent cotton placed in the apex of a sterile funnel.

(e) The washings should be adjusted (using 12 percent sodium chloride containing 0.25 to 0.5 percent phenol) so that the turbidity is 50 times greater than tube 0.75 of McFarland's nephelometer, or to a reading of 7 mm. by the Gates nephelometer.

(f) The individual strain antigens should be tested with negative sera for their insensitivity and with positive sera for high agglutinability in comparison with known satisfactory antigen. The antigens of the separate strains should be combined in equal volume-density and stored in the refrigerator (5° to 10°C) in tightly stoppered bottles.

(g) The tests should be conducted on a suitable, smooth plate. The serum-antigen dilution should be made so that the dilution will not exceed 1:50 when compared to the standard tube agglutination test. When testing turkey blood samples, it is desirable to use a serum-antigen dilution equivalent to the 1:25 in the tube method. The serum should be added to the antigen and mixed thoroughly by use of the tip of the serum pipette. Most strong positive reactions will be plainly evident within 15 to 20 seconds. The final reading should be made at the end of 2 or 3 minutes. Heating the plate at approximately 37°C will hasten agglutination. Before reading, the plate should be rotated several times.

(h) The results shall be recorded as described in §147.1(h).

§147.3 The stained antigen, rapid whole-blood test.[3/]

(a) The description of the preparation of antigen is not herein included because the antigen is a proprietary product produced only under license from the Secretary of Agriculture.

(b) A loop for measuring the correct quantity of blood can usually be obtained from the manufacturer of the antigen. A satisfactory loop may be made from a piece of No. 20 gage nichrome wire, 2 1/2 inches long, at the end of which is fashioned a loop three-sixteenths of an inch in diameter. Such a loop, when filled with blood so that the blood appears to bulge, delivers 0.02 cc. A medicine dropper whose tip is adjusted to deliver 0.05 cc. is used to measure the antigen. A glass plate about 15 inches square, providing space for 48 tests, has proved satisfactory for this work. The use of such a plate enables the tester to have a number of successive test mixtures under observation without holding up the work to wait for results before proceeding to the next bird.

3/ The procedure described is a modification of the method reported by Schaffer, MacDonald, Hall, and Bunyea, Jour. Amer. Vet. Med. Assoc. 79 (N. S. 32): 236-240 (1931).

(c) A drop of antigen should be placed on the testing plate. A loopful of blood should be taken up from the wing vein. When submerged in the blood and then carefully withdrawn, the loop becomes properly filled. On looking down edgewise at the filled loop, one observes that the blood appears to bulge. The loopful of blood then should be stirred into the drop of antigen, and the mixture spread to a diameter of about 1 inch. The loop then should be rinsed in clean water and dried by touching it to a piece of clean blotting paper, if necessary. The test plate should be rocked from side to side a few times to mix the antigen and blood thoroughly, and to facilitate agglutination. The antigen should be used according to the directions of the producer.

(d) Various degrees of reaction are observed in this as in other agglutination tests. The greater the agglutination ability of the blood, the more rapid the clumping and the larger the clumps. A positive reaction consists of a definite clumping of the antigen surrounded by clear spaces. Such reaction is easily distinguished against a white background. A somewhat weaker reaction consists of small but still clearly visible clumps of antigen surrounded by spaces only partially clear. Between this point and a negative or homogeneous smear, there sometimes occurs a very fine granulation barely visible to the naked eye; this should be disregarded in making a diagnosis. The very fine marginal clumping which may occur just before drying up is also regarded as negative. In a nonreactor, the smear remains homogeneous. (Allowance should be made for differences in the sensitiveness of different antigens and different set-ups, and therefore, a certain amount of independent, intelligent judgment must be exercised at all times. Also, the histories of the flocks require consideration. In flocks where individuals show a suspicious agglutination, it is desirable to examine representative birds bacteriologically to determine the presence or absence of S. pullorum.)

§147.4 The tube agglutination test for S. typhimurium.

(a) The procedure for the collection and delivery of blood samples in the tube agglutination test for S. typhimurium is the same as that described in §147.1(a).

(b) The "O" antigen should be prepared as follows:

(b)(1) The antigen shall consist of a representative nonmotile strain of S. typhimurium which is of known antigenic composition and high agglutinability but is not sensitive to negative and nonspecific sera. Strain P 10 meets these requirements.

(b)(2) The stock culture is maintained on 1 percent nutrient agar deeps, which have been incubated for 18-24 hours at 37°C. They are stored at room temperature.

(b)(3) A satisfactory medium used for growing the organism is veal infusion agar (Difco). It is dispensed in 50 ml amounts into 500 ml medicine bottles, with screw caps, and sterilized at 15 pounds pressure for 20 minutes. The bottles are then laid flat upon an even surface until the medium has solidified.

(b)(4) The inoculum used for preparation of "O" antigen is a nonmotile strain of S. typhimurium. The organism is grown in veal infusion broth (Difco) for 18-24 hours at 37°C; then plated, for single colony isolation, on veal infusion agar plates. These plates are incubated for 18-24 hours at 37°C. After incubation, single colonies are picked and transferred to veal infusion agar slants, which are incubated for 18-24 hours at 37°C. After this, the cultures are tested for smoothness by using a 1:500 dilution of acriflavine.

(b)(5) Smooth cultures are inoculated into flasks containing veal or beef infusion broth which is incubated for 18-24 hours at 37°C. The incubated broth suspension of organisms is dispensed into the antigen bottles containing veal infusion agar. The suspension is distributed evenly over the agar surface by gently tilting the bottles from side to side. The inoculated bottles are then laid flat, agar side down, for 10-20 minutes. They are subsequently incubated, agar side upward, for 24-48 hours at 37°C before harvesting.

(b)(6) The harvesting of the organism consists of washing the growth from each antigen bottle with 0.5 percent phenolized physiological saline. The bacterial suspension from each bottle is filtered through sterile milk pad filters into a large sterile container or through a thin layer of absorbent cotton in a Buchner funnel with the aid of suction. To each 100 ml. of the bacterial suspension is added additional phenol to make the final concentration 0.5 percent. The concentrated antigen is tested for sterility at intervals after 24 hours. After sterility is proved, the stock antigen is standardized to determine the density according to the McFarland nephelometer scale.

(b)(7) The diluted antigen to be used in routine testing is prepared from stock antigen, by diluting with 0.25 percent phenolized saline, and is standardized to a turbidity corresponding to 0.75-1.00 of the McFarland nephelometer scale.

(c) The maximum serum dilution employed for the "O" antigen tube test must not exceed 1:25. In all official reports on the blood test, the serum dilutions should be indicated. The sera should be introduced into the agglutination tubes in the desired amounts with well-cleaned serological pipettes or special serum delivery devices which do not permit the mixing of different sera. The antigen and serum should be well mixed before incubation. The serum and antigen mixture must be incubated for at least 20 hours at 37°C.

(d) The results shall be recorded as described in §147.1(h).

§147.5 The microagglutination test for pullorum-typhoid.

Routinely, the microagglutination test is applied as a single-dilution test and only a single 18-24 hour reading is made.

(a) The procedure for the collection and delivery of blood samples in the microagglutination test is the same as that described in §147.1(a). A method that has proven advantageous is to transfer the serum samples from the blood clot to a microplate as described in "Applied Microbiology," volume 24, No. 4, October 1972, pages 671-672. The dilutions are then performed according to paragraphs (d) or (e) of this section.

(b) Stained microtest antigen for pullorum-typhoid is supplied as concentrated stock suspension and must be approved by the Department.[4/] Direction for diluting will be provided with the antigen. The stock as well as the diluted antigen prepared each day should be kept sealed in the dark at 5° to 10°C when not in use.

[4/] Information as to criteria and procedures for approval of concentrated stock suspension of stained microtest antigens may be obtained from the National Poultry Improvement Plan Staff, Building 265, Beltsville Agricultural Research Center-East, Beltsville, Maryland 20705.

(c) Available data indicate that a 1:20 dilution for the microagglutination test is most efficient for the detection of pullorum-typhoid agglutinins in both chickens and turkeys. In all official reports on the blood test, the serum dilutions shall be indicated.

(d) The recommended procedure for the 1:20 dilution in the microagglutination test is as follows:

(d)(1) Add 100 microliters (0.10 cc) of 0.85 percent physiological saline to each well of the microplate.

(d)(2) Using a microdiluter or a multimicrodiluter handle fitted with twelve 10 microliter microdiluters, transfer 10 microliters (0.01 cc) of the serum sample from the collected specimen to the corresponding well of the microplate. This is accomplished by touching the surface of the serum sample with the microdiluter and then transferring and mixing with the diluent in the microplate well. The microdiluter is removed, blotted, touched to the surface of the distilled water wash, and again blotted. Other acceptable methods of serum delivery are described in "Applied Microbiology," volume 21, No. 3, March 1971, pages 394-399.

(d)(3) Dilute the microtest antigens with 0.50 percent phenolized saline and add 100 microliters (0.1 cc) to each microplate well.

(d)(4) Seal each plate with a plastic sealer or place unsealed in a tight incubation box as described in "Applied Microbiology," volume 23, No. 5, May 1972, pages 931-937. Incubate at 37°C for 18-24 hours.

(d)(5) Read the test results as described in paragraph (f) of this section.

(e) The recommended procedure for a microagglutination test titration is as follows:

(e)(1) Add 50 microliters (0.05 cc) of 0.85 percent physiological saline to each well of the microplate.

(e)(2) To the wells representative of the lowest dilution in the titration, add an additional 50 microliters (0.05 cc) of 0.85 percent physological saline making a total of 100 microliters in these wells.

(e)(3) Transfer each serum sample as described in §147.5(d)(2) of this section to the first well containing 100 microliters (0.10 cc) in the titration, which represents the lowest dilution.

(e)(4) Make two-fold serial dilutions of each serum by transferring 50 microliters (0.05 cc) of diluted serum from one well to the next using twelve 50 microliter microdiluters fitted in a multimicrodiluter handle. When transfers have been made to all of the wells of the desired series, the 50 microliters remaining in the microdiluters are removed by blotting, touching the microdiluters to the surface of the distilled water wash, and blotting again.

(e)(5) Dilute the desired microtest antigen with 0.50 percent phenolized saline and add 50 microliters (0.05 cc) to each microplate well.

(e)(6) Seal each plate with a plastic sealer or place the unsealed microplates in a tight incubation box and incubate at 37°C for 18-24 hours.

(e)(7) Read the test results as described in paragraph (f) of this section.

(f) Read the test results with the aid of a reading mirror. Results are interpreted as follows:

(f)(1) N, or - (negative) when the microplate well has a large, distinct button of stained cells; or

(f)(2) P, or + (positive) when the microplate well reveals no antigen button; or

(f)(3) S, or ? (suspicious) when the microplate well has a small button. Suspicious reactions may tend to be more positive than negative [+] or vice versa [∓] and can be so noted if desired.

§147.6 Procedure for determining the status of flocks reacting to tests for Mycoplasma gallisepticum and Mycoplasma synoviae.

The macroagglutination tests for Mycoplasma antibodies, as described in "Standard Methods for Testing Avian Sera for the Presence of Mycoplasma Gallisepticum Antibodies" published by the Agricultural Research Service, USDA, March 1966, and the microagglutination tests, as reported in the Proceedings, Sixteenth Annual Meeting of the American Association of Veterinary Laboratory Diagnosticians, 1973, shall be the official tests. Procedures for isolation and identification of Mycoplasma may be found in Isolation and Identification of Avian Pathogens, published by the American Association of Avian Pathologists and §§147.15 and 147.16 of this part.

(a) When reactors are submitted to a laboratory as prescribed by the Official State Agency, the following criteria shall be used to determine if the flock is positive for M. gallisepticum or M. synoviae:

(a)(1) Active air sac lesions, sinusitis, synovitis, or other clinical signs of a respiratory disease;

(a)(2) Recovery by culture of the Mycoplasma for which the flock was tested;

(a)(3) Supplemental serological test.

(b) If all of these tests are negative, the flock shall be deemed to have had no reactors for the Mycoplasma for which the flock was tested. If the Mycoplasma for which the flock was tested is isolated bacteriologically, the flock shall be considered infected. If any of the other tests described in paragraphs (a)(1) or (3) of this section is positive, the flock shall be considered suspicious, and additional agglutination and hemagglutination inhibition (HI) tests shall be conducted according to the following sequence:

(b)(1) If the tube agglutination or the serum plate test is negative, the flock qualifies.

(b)(2) If the tube agglutination or the serum plate test is positive, the hemagglutination inhibition (HI) test and/or the serum plate dilution (SPD) test shall be conducted.

(b)(3) If the tube agglutination or serum plate tests are positive and HI and/or the SPD tests are negative, the flock shall be retested in accordance with paragraph (b)(6) of this section.

(b)(4) If HI titers of 1:40 or SPD titers of 1:5 are found, the flock shall be considered suspicious and shall be retested in accordance with paragraph (b)(6) of this section.

(b)(5) If HI titers of 1:80 or SPD titers of 1:10 or higher are found, the flock shall be considered infected: Provided, That, at the discretion of the Official State Agency, additional tests may be conducted in accordance with paragraph (b)(6) of this section before final determination of the flock status is made.

(b)(6) Fourteen days after the previous bleeding date, all birds or a random sample comprised of 75 birds shall be tested by the serum plate or tube agglutination test. Tested birds shall be identified by numbered bands.

(b)(7) If the tube agglutination test or serum plate test is negative for the Mycoplasma for which the flock was tested, the flock qualifies.

(b)(8) If the tube agglutination or serum plate test is positive, the HI and/or SPD test shall be conducted on the reacting samples.

(b)(9) On the retest, if the tube agglutination or serum plate tests are positive at the same or higher rate and the HI or SPD tests are negative, the flock shall be considered suspicious and shall be retested in accordance with paragraph (b)(6) of this section.

(b)(10) On the retest if HI titers of 1:80 and/or SPD titers of 1:10 or higher are found, the flock shall be considered infected: Provided, That, at the discretion of the Official State Agency, additional tests may be conducted in accordance with paragraph (b)(6) of this section before final determination of the flock status is made.

(b)(11) If HI titers of 1:80 and/or SPD titers of 1:10 or higher are found on the second retest, the flock shall be considered infected for the Mycoplasma for which it was tested.

(b)(12) If the tube agglutination or serum plate tests are found on the second retest to be positive at the same or higher rate and the HI and/or SPD tests are negative, the flock should be considered infected: Provided, That if the status of the flock is considered to be equivocal, the Official State Agency may examine reactors by the in vivo bio-assay and/or culture procedures before final determination of the flock status is made.

(b)(13) If the in vivo bio-assay and culture procedures are both negative, the Official State Agency may qualify the flock for the classification for which it was tested.

(b)(14) If the in vivo bio-assay or culture procedures are positive, the flock shall be considered infected: Provided, That if only the bio-assay is positive, additional in vivo bio-assay or cultural examinations may be conducted by the Official State Agency before final determination of the flock status is made.

(b)(15) If the in vivo bio-assay or cultures are positive on retest, the flock shall be considered infected for the mycoplasma for which it was tested.

Subpart B - Bacteriological Examination Procedures

§147.11 Laboratory procedure recommended for the bacteriological examination of salmonella reactors.

(a) Grossly normal or diseased liver, heart, pericardial sac, spleen, lung, kidney, pancreas, peritoneum, drained gallbladder, oviduct, misshapen ova, testes, inflamed or unabsorbed yolk sac, and other visibly pathological tissues where purulent, necrotic, or proliferative lesions are seen (including cysts, abscesses, hypopyon, and inflamed serosal surfaces), should be directly cultured by means of a flamed wire loop or with sterile swabs.[1/] Careful aseptic technique must be utilized throughout the process of collecting tissues. Selective media should not be relied upon to deal with contaminants, since some strains may not dependably survive and grow in certain selective media. Inoculate veal infusion (VI) and brilliant green (BG) agar plates. Incubate the plates for 24 and 48 hours at 37°C. The digestive system should always be cultured spearately (see paragraph (f) of this section) after other anatomical organs and systems have been collected and cultured.

1/ Culture media preparation and biochemical identification charts can be obtained from Culture Methods for the Detection of Animal Salmonellosis and Arizonosis, Committee on Salmonellosis and Arizonosis, AAVLD, 1976 Iowa State University Press, Ames, Iowa 50010.

(b) The following organs should be aseptically collected for culture:

(b)(1) Heart (apex, pericardial sac, and contents if present.);

(b)(2) Liver (portions exhibiting lesions or, in grossly normal organs, the drained gallbladder and adjacent liver tissues.);

(b)(3) Ovary-Testes (entire inactive ovary or testes, but if ovary is active, use own judgment and include any atypical ova.);

(b)(4) Oviduct (if active, include any debris and dehydrated ova.);

(b)(5) Pancreas and kidneys; and

(b)(6) Spleen.

(c) Aseptically collect 10-15 g or whatever lesser amount is available of each organ or site listed in paragraph (b) from each reactor, and grind or blend them completely in 10 times their volume of VI broth. Organs may be processed individually or in combinations where appropriate. Suspensions should be transferred in 10-ml aliquots to 100 ml of both VI and tetrathionate brilliant green (TBG) broth and incubated at 37° for 24 hours. Plate the VI broth on VI and BG agar and plate the TBG broth on BG agar and incubate at 37°. Examine these plates after 24 and 48 hours of incubation. The contents of the gallbladder can be cultured separately by inoculating 10 ml of VI and TBG broth with cotton swabs and incubating at 37°C for 24 hours. Plate on BG agar and incubate at 37°. Examine these plates after 24 and 48 hours of incubation. If contamination with pseudomonas or proteus is a problem, make platings on BG sulfapyridine (BGS) agar.

(d) Where field samples are directly inoculated into enrichment broths and a delay of several days occurs before they reach a laboratory, or if recovery of low numbers or organisms is expected from a primary culture, a secondary enrichment culture should be prepared. Subculture a week-old primary culture by transferring 1 ml of inoculum into a fresh tube 10 ml of enrichment broth. This seconary enrichment should be incubated at 37°C for 24 hours before plating. (See paragraph (a) of this section.) TBG broth is recommended for this procedure.

(e) Make a composite sample of the following parts of grossly normal or diseased tissues from the digestive tract: Crop wall, duodenum, jejunum (including remnant of yolk-sac attachment), both ceca, cecal tonsils, and rectum-cloaca. Aseptically collect 10-15 g or whatever lesser amount is available of each organ or tissue and grind or blend them completely in 10 times their volume of TBG broth. Transfer 10 ml of a composite sample of a suspension from the digestive tract into 100 ml of TBG broth, and incubate flasks at 42°C for 24 hours. Cultures may be incubated at 37°C if 42°C incubators are not available. The higher incubation temperatures for TBG broth reduce populations of competitive contaminants common in gut tissue. Plate on BG agar and incubate at 37°C. Examine the plates after 24 and 48 hours of incubation. If contamination with pseudomonas or proteus is a problem make plating on BGS agar.

(f) If preferred, individual cotton swab samples may also be taken from the upper, middle, and lower intestinal tract (including both ceca and the rectum-cloaca). Deposit swabs in 10 ml of TBG broth and incubate and plate as described in paragraph (e) of this section.

(g) Transfer suspect colonies to triple-sugar-iron (TSI) agar and lysine-iron (LI) agar and incubate at 37°C for 24 hours.

(h) Cultures revealing typical reactions of salmonellae or arizonae on TSI and LI agar slants should be transferred to any of the following appropriate biochemical tests for final identification: Dextrose, lactose, sucrose, mannitol, maltose, dulcitol, malonate, gelatin, urea broth, citrate, lysine decarboxylase, orinthine decarboxylase, methyl red and Voges-Proskauer, KCN, salicin broths, indole, and hydrogen sulfide.

Motility or non-motility is demonstrated by inoculating a suitable semisolid medium.[2/] The Analytical Profile Index for Enterobacteriacae (API) system may be utilized for identification if feasible. For arizonae identification, make readings daily up to 10 days. An O-nitrophenyl-beta-D-galactopyransíde (ONPG) disc may be used to identify slow lactose fermeters.[3/]

(i) All salmonella cultures should be serologically typed.

§147.12 Procedures for collecting environmental samples and cloacal swabs for bacteriological examination.

Information concerning the pen arrangement and number of birds per pen should be obtained from the owner so that the required number of samples per pen and per flock can be determined. A means of identifying each sample by pen of origin should be provided. The vehicle transporting the personnel taking the samples should be left as far as practical from the poultry pens. Sanitary precautions, including personal cleanliness, should be observed during the sampling procedure. The hands should be carefully washed with a sanitizing soap prior to the sampling. Outer clothing, including gloves, should be changed between visits to different premises so that clean clothing is worn upon entering each premises.

The used and clean apparel should be kept separate. Boots or footwear should be cleaned and disinfected between visits to different premises. Disposable caps should be provided and discarded after use on each premises. After collection, the samples should be protected from drying, light, and excessive temperatures and delivered to the laboratory within one day. If delivery is delayed, samples should be refrigerated.

(a) Environmental samples. Fecal material, litter, or dust to be submitted for bacteriological examination should be collected in accordance with the procedures described in paragraphs (a)(1) or (2) of this section:

(a)(1) Procedure for sampling in broth. Authorized laboratories will provide capped tubes 1-2 cm in diameter and 15-20 cm in length which are two-thirds full of a recently made, refrigerated, sterile enrichment broth (Selenite Brilliant Green Sulfapyridine or Tetrathionate Brilliant Green) for each sample. Sufficient tubes should be taken to the premises to provide at least one tube per pen or one tube per 500 birds, whichever is greater. At least one sterile, cotton-tipped applicator will be needed for each tube. The dry applicator is first placed or drawn through fresh manure (under roost, near water troughs, cecal droppings, or diarrhetic droppings). After this and each subsequent streaking, the cotton-tipped applicator is placed in the tube of broth and swirled to remove the collected material. The applicator is then withdrawn and is used for taking additional specimens by streaking on or through areas where defecation, trampling of feces, or settling of dust are common; i.e., on or near waterers, feeders, nests, or rafters, etc.

2/ Formulation for the semisolid motility medium can be obtained from: Isolation and Identification of Avian Pathogens, American Association of Avian Pathologist, Texas A & M University, College Station, Texas 77843, 1980.

3/ ONPG discs are available from: Baltimore Biological Laboratories, Cockeysville, Maryland 21030.

When the volume of material collected equals approximately 10 percent of the volume of the broth (usually 10-12 streakings), the applicator is placed in the tube and the stick is broken in half.
The lower or cotton-tipped half is left in the broth, and the upper half is retained for future disposal. The cap is then replaced on the inoculated tube, and the sampling procedure is continued in other areas of the pen.

(a)(2) Procedure for sampling in dry containers. A sample of fecal material, litter, or dust is placed in a sterile, sealable container. The sample shall consist of several specimens of material taken from a representative location in the pen. At least 10 g (approximately a heaping tablespoonful) of material shall be collected for each sample. The specimens in each sample shall be collected with a sterile tongue depressor or similar uncontaminated instrument. The samples should vary in type and consistency. Half of the samples should be comprised of material representing defecated matter from a large portion of the flock; i.e., trampled, caked material near waterers and feeders. The minimum number of samples to be taken shall be determined by the following:

Five samples from pens of up to 500 birds;
Ten samples from pens of 500 to 2,500 birds;
Fifteen samples from pens with more than 2,500 birds.

(b) Cloacal swabs. Cloacal swabs for bacteriological examination are taken from each bird in the flock or from a minimum of 500 birds in accordance with the procedure described in paragraph (a)(1) of this section.

(b)(1) Procedure for taking cloacal swabs. The authorized laboratory will provide sterile capped tubes or other suitable containers and cotton-tipped applicators for use in taking the cloacal swabs. The cotton-tipped applicator is inserted into the cloaca and rectum in such a manner as to insure the collection of fecal material. The swab and adhering fecal material is then placed in the tube and the stick is broken in half, with the upper half retained for future disposal. The cloacal swabs may be combined in the sterile tubes in multiples of five or in combinations specified by the authorized laboratory.

§147.13 Procedure for bacteriological culturing of eggshells for for colon bacilli organisms.

Proper precautions to avoid environmental contamination of the samples during the collection and laboratory process, and proper handling of the samples following collection are essential. Each State Inspector involved in eggshell culture activities must receive instruction in the necessary sanitation procedures, sampling procedures, and sample handling by the authorized laboratory involved. The Official State Agency will maintain a record showing that the required instruction was given to each State Inspector.

(a) Sample selection. Forty (40) eggs in the top flats of each of three randomly selected cases of sanitized eggs from each flock will be utilized for each sampling.

(b) Swab procedure. A 2.5 centimeter diameter circular area of the large end of each of the eggs will be rubbed with a sterile swab previously moistened with sterile lactose broth, or other suitable liquid media provided by the authorized laboratory. One swab will be used for five eggs, and four swabs will be pooled to each sterile, capped tube provided by the authorized laboratory.

(b)(1) From the tube containing four swabs and lactose broth or other suitable media, 1 ml will be transferred to 10 ml lactose in a fermentation tube.

(b)(2) Incubate at 37°C for 48 hours. The presence of acid, and gas in the amount of 10 percent or more after 24 and 48 hours of incubation, provides a presumptive conclusion of the presence of colon bacilli organisms.

§147.14 Procedures to determine status and effectiveness of sanitation monitored program.[4/]

The following monitoring procedures may be applied at the discretion of the Official State Agency:

(a) Monitor effectiveness of sanitation program.

(a)(1) Culture the surface of cased eggs periodically for fecal contaminating organisms as described in §147.13.

(a)(2) Culture a sample of dead-in-shell eggs periodically from each breeding flock for coliforms. The culture media will be designed to include detection of salmonella species.

§147.15 Laboratory procedure recommended for the bacterioloical examination of mycoplasma reactors.[5/]

(a) Turbinates, trachea, air sacs, sinuses, nasal passages, respiratory exudates, synovial fluid, eggs (including yolk, yolk sacs, membranes and allantoic fluid), should be directly sampled with sterile swabs. Aseptic techniques are very important as some organisms may not be suppressed by the antimicrobial agents used in this procedure. Tissue suspensions from large volumes are sometimes desirable from the sites listed above and occasionally from the oviduct and cloaca. Tissues should be ground or blended completely in 10 times their volume of Mycoplasma Broth Medium (MBM). (See paragraph (e) of this section.) Specimens submitted to referral laboratories in order of preference for recovery of the mycoplasma organisms are: (1) live birds, (2) refrigerated fresh tissues, (3) tissue specimens packed with dry ice.

(b) Inoculate 5-10 ml of MBM with a swab, wire loop or 0.1 ml of the tissue suspension. When evidence of growth is observed (lowered pH or turbidity of broth) transfer each broth culture as needed to maintain the original isolates. Incubate tubes at 37°C for at least 21 days before discarding as negative.

4/ Laboratory procedures for monitoring operations proposed here are described in the following two publications: (a) Isolation and Identification of Avian Pathogens, American Association of Avian Pathologists, Texas A & M University, College Station, Texas 77843, 1975 and (b) Culture Methods for the Detection of Animal Salmonellosis and Arizonosis, Iowa State University Press, Ames, Iowa 50010, 1976.

5/ Yoder, H. W., Jr., "Mycoplasmosis." In: Isolation and Identification of Avian Pathogens. (Stephen B. Hitchner, Chairman, Charles H. Domermuth, H. Graham Purchase, James E. Williams.) 1980, pp. 40-42, Creative Printing Company, Inc., Endwell, New York 13760.

When growth is first observed or if no growth occurs by the 4th or 5th day of incubation, inoculate broth culture onto a plate of Mycoplasma Agar Medium (MAM). (See paragraph (f) of this section.) Several cultures may be inoculated on one plate by using a wire loop or a cotton swab. Incubate plates 3-5 days at 37°C in a high humidity-chamber. If preferred, 5 percent CO_2 may be added or a candle jar may be used. Tiny circular and translucent colonies with elevated centers are very suggestive of mycoplasma. Indirect lighting and a low power or dissecting microscope are recommended for observation of the colonies as they are rarely more than 0.2-0.3 mm in diameter.

(c) Isolates must be serotyped.

(c)(1) Isolates may be shipped in MBM with ice packs if shipment will be in transit less than 2-3 days. Longer shipments require freezing of the MBM with dry ice, or shipping MAM slants at room temperature. Isolates must have indications of growth before shipment is made.

(c)(2) Isolates may be stored in MBM at -20°C for 2-3 weeks, or they may be stored at -68°C for several years.

(d) Alternate method of culture: An overlay enrichment culture for fastidious and sensitive mycoplasma, especially for M. meleagridis should be included.

(d)(1) Pour 2-3 ml of MAM into a test tube and tilt the tube until a slant (approximately 45°) is obtained. Other containers are acceptable.

(d)(2) Overlay the slant with sufficient MBM, so that the media (including inoculum) covers the agar slope.

(d)(3) Inoculate the culture as indicated in paragraph (b) of this section.

(d)(4) Incubate and examine the overlay as indicated in paragraph (b) of this section.

(e) Preparation of media components:[6/]

(e)(1) Deionized distilled water suitable for cell culture fluids should be used.

(e)(2) All glassware should be carefully washed with a non-residue detergent such as Alcojet[7/] and rinsed three times in tap water and twice in deionized distilled water.

(e)(3) Thallium acetate in a 10 percent solution is added to an approximate final concentration of 1:4000; however, highly contaminated specimens may require a final concentration of 1:2000.[8/] Thallium acetate is added to deionized distilled water first, except as noted in paragraph (e)(4) of this section, to prevent the precipitation of proteins.

6/ Trade names are used in these procedures solely for the purpose of providing specific information. Mention of a trade name does not constitute a guarantee or warranty of the product by the U.S. Department of Agricultue or an endorsement over other products not mentioned.

7/ Alcojet is available from: Alconox, Inc., New York, New York 10003.

8/ Thallium acetate may be obtained from Fischer Scientific Company.

(e)(4) Mycoplasma Broth Base, dextrose, phenol red, and cysteine hydrochloride are added to deionized distilled water first if autoclave sterilization is used.[9] Thallium acetate and then the remaining components are added aseptically after cooling the autoclaved media to 45°C or less.

(e)(5) Use sterile deionized distilled water to reconstitute penicillin.

(e)(6) Sterile serum should be inactivated by heating at 56°C for 30 minutes. Swine serum may be used for M. gallisepticum, M. synoviae, M. gallinarum, and M. meleagridis isolation; however, horse serum is usually recommended for M. meleagridis isolation.

(e)(7) Phenol red should be prepared as a 1 percent solution.

(e)(8) NAD (beta nicotinamide adenine dinucleotide or coenzyme I) should be prepared as a 1 percent solution.[10]

(e)(9) Cysteine hydrochloride, prepared as a 1 percent solution, is used to reduce the NAD for M. synoviae growth.

(e)(10) A purified agar product such as Nobel (Special agar) is used in the MAM.[11]

(e)(11) Adjust the pH with NaOH.

(e)(12) Sterilization may be accomplished by two methods:

(e)(12)(i) Filtration sterilization through a 0.20 micron filter is the recommended method. Aseptic techniques must be utilized.

(e)(12)(ii) Autoclave sterilization at 120°C, 15 pounds pressure (103 kPa), for 15 minutes may be used, if preferred, when following the procedure described in paragraph (e)(4) of this section.

(e)(13) Phenol red, dextrose, and NAD may be omitted when culturing for M. meleagridis and M. gallinarum.

(e)(14) When culturing for M. meleagridis from contaminated samples include 100 units/ml of Polymyxin B in MBM.

(f) Mycoplasam Broth Medium (Frey) is prepared as follows: To 850-880 ml of deionized distilled water;

Add:

Thallium acetate (ml)	2.5	(1:4000)
Potentially contaminated samples (ml)	5.0	(1:2000)
Mycoplasma Broth Base (g)	22.5	
Aqueous penicillin (units)	500,000	
Sterile serum (ml)	120 to 150.0	
Phenol red plus (ml)	2.5	
NAD (ml)	12.5	
Cysteine hydrochloride (ml)	12.5	
Dextrose (g)	1.0-1.5	
Adjust pH to 7.8		
Filter sterilize		

9/ Mycoplasma Broth Base may be obtained from: (a) Product #M 33600, Gibco Diagnostics, 2801 Industrail Drive, Madison, Wisconsin 53711. (b) Product #3900-3212, Scott Laboratories, Inc., 8 Westchester Plaza, Elmsford, New York 10523.

10/ NAD Grade III may be obtained from: Sigma Chemical Company, P.O. Box 14508, St. Louis, Missouri 63178.

11/ Nobel Agar may be obtained from: Difco Laboratories, Box 1058-A, Detroit, Michigan 48201.

(f)(1) Broth may be stored at 4°C for at least 2 weeks or at -40°C for longer periods.

(g) Mycoplasma Agar Medium (Frey) is prepared as follows:
to 850-880 ml of deionized distilled water;

Add:

Mycoplasma Broth Base (g)	22.5	
Adjust pH to 7.8		
Purified agar (g)	18.0	
Autoclave and cool in 45°C water bath		
Thallium acetate (ml)	2.0	(1:4000)
Sterile serum at 45°C (ml)	150.0	
Aqueous penicillin (units)	400,000	
NAD (ml)	12.5	
Cysteine hydrochloride (ml)	12.5	

(g)(1) Rotate flask gently and pour about 15 ml of media into each petri dish.

(g)(2) Stack petri dishes only 2-3 high in a 37°C incubator up to 2 hours to remove excess moisture.

(g)(3) Wrap inverted plates in sealed bundles and store at 4°C for not more than 15 days.

(h) New component or media batches should be monitored to compensate for changes in formulation due to alterations of purity, concentration, preparation, etc. A known series of titrations from a single culture should be made on both new and old media. The media should be compared on the basis of growth, colony size, and numbers of colonies which develop.[12/]

§147.16 Procedure for the evaluation of mycoplasma reactors by in vivo bio-assay (enrichment).

This procedure has been shown to be sensitive enough to detect less than 100 mycoplasma organisms under proper conditions.[13/] Proper conditions are defined in this section.

(a) Obtain chickens or turkeys (test birds) which are at least 3 weeks of age and are free of *M. gallisepticum*, *M. synoviae*, and *M. meleagridis* and transport them in a manner to prevent their being contaminated by any infectious avian disease.

(a)(1) Maintain test birds in an area that has been effectively cleaned and disinfected.

(a)(2) The area should be isolated from other birds or animals.

12/ "Laboratory Procedures and Medium For The Isolation Of Mycoplasma From Clinical Materials." Laboratory Diagnosis of Mycoplasma in Food Animals, Proceedings of Nineteenth Annual Meeting, The American Association of Veterinary Laboratory Diagnosticians, 1976, pp. 106-115, AAVLD, 6101 Mineral Point Road, Madison, Wisconsin 53705.

13/ Research results are described in the following two publications: (a) Bigland, C. H. and A. J. DaMassa, "A Bio-Assay for Mycoplasma Gallisepticum." In: United States Livestock Sanitary Association Proceedings, 67th, 1963, pp. 541-549. (b) McMartin, D. A., "Mycoplasma Gallisepticum in the Respiratory Tract of the Fowl." In: The Veterinary Record, September 23, 1967, pp. 317-320.

(a)(3) Personnel caring for the test birds should take the necessary precautions (see §147.26(b)) to prevent the mechanical transfer of infectious avian diseases from other sources.

(b) Test birds to be used for inoculation with contaminated tissues should be serologically negative by the serum plate agglutination test.

(b)(1) Inoculated test birds should be isolated from non-inoculated control birds for the length of any experiment.

(c) Aseptically obtain tracheal, turbinate, and sinus mucosa, lung and sinus exudates, cervical, thoracic, and abdominal air sac tissues (including lesions), portions of the oviduct, and synovial fluid from at least four suspect, donor birds. In a sterile device, blend the tissues completely in four times their volume of Mycoplasma Broth Medium (Frey), (see §147.15(e)). Suspensions may be made from tissue pools. Inoculate test birds within 30 minutes of preparation of suspensions.

(c)(1) Inoculate at least four test birds for each suspension pool via the abdominal air sac and infraorbital sinus, with up to 1/2 ml of inoculum per site.

(c)(2) Test birds should be bled every 7 days for 35 days to identify sero-converters.

(c)(3) At 35 days, test birds should be sacrificed and bacteriologic isolation and identification of mycoplasma attempted (see §147.15). Note especially the sites of inoculation for typical gross or microscopic mycoplasma lesions.

(d) Donor birds are considered infected when:

(d)(1) Test birds have serum plate antibodies for the mycoplasma for which the donor birds were tested, regardless of HI test results, and control birds stay serologically negative; or

(d)(2) Mycoplasma organisms are isolated from the test birds and serotyped positive for the mycoplasma for which the donor birds were tested, and control birds stay serologically and culturally negative.

(e) Laboratory findings may be verified by direct culture of material from sick birds or by inoculating seronegative birds from the suspect flock and comparing serological findings with those from the test birds.

Subpart C - Sanitation Procedures

§147.21 Flock sanitation.

To aid in the maintenance of healthy flocks, the following procedures should be practiced:

(a) Baby poultry should be started in a clean brooder house and maintained in constant isolation from older birds and other animals. Personnel that are in contact with older birds and other animals should take precautions, including disinfection of footwear and change of outer clothing, to prevent the introduction of infection through droppings that may adhere to the shoes, clothing, or hands. (See §147.24(a).)

(b) Range used for growing young stock should not have been used for poultry the preceding year. Where broods of different ages must be kept on the same farm, there should be complete depopulation of brooder houses and other premises following infection of such premises by any contagious disease.

(c) Poultry houses should be screened and proofed against free-flying birds. An active rodent eradication campaign is an essential part of the general sanitation program. The area adjacent to the poultry house should be kept free from accumulated manure, rubbish, and unnecessary equipment. Dogs, cats, sheep, cattle, horses, and swine should never have access to poultry operations. Visitors should not be admitted to poultry areas, and authorized personnel should take the necessary precautions to prevent the introduction of disease.

(d) Poultry houses and equipment should be thoroughly cleaned and disinfected prior to use for a new lot of birds. (See §147.24(a).) Feed and water containers should be situated where they connot be contaminated by droppings and should be frequently cleaned and disinfected. Dropping boards or pits should be constructed so birds do not have access to the droppings.

(e) Replacement breeders shall be housed at the proper density consistent with the type of building and locality and which will allow the litter to be maintained in a dry condition. Frequent stirring of the litter may be necessary to reduce excess moisture and prevent surface accumulation of droppings. Slat or wire floors should be constructed so as to permit free passage of droppings and to prevent the birds from coming in contact with the droppings. Nesting areas should be kept clean and, where appropriate, filled with clean nesting material.

(f) When an outbreak of disease occurs in a flock, dead or sick birds should be taken, by private carrier, to a diagnostic laboratory for complete examination. All salmonella and arizona cultures isolated should be typed serologically, and complete records maintained by the laboratory as to types recovered from each flock within an area. Records on isolations and serological types should be made available to Official State Agencies or other animal disease control regulatory agencies in the respective States for followup of foci of infection. Such information is necessary for the development of an effective salmonella control program.

(g) Introduction of started or mature birds should be avoided to reduce the possible hazard of introducing infectious diseases. If birds are to be introduced, the health status of both the flock and introduced birds should be evaluated.

(h) In rearing broiler or replacement stock, a sound and adequate immunization program should be adopted. Since different geographic areas may require certain specific recommendations, the program recommended by the State experiment station or other State agencies should be followed.

(i) Feed, pelleted by heat process, should be fed to all age groups. Proper feed pelleting procedures can destroy many disease producing organisms contaminating feedstuffs.

§147.22 Hatching egg sanitation.

Hatching eggs should be collected from the nests at frequent intervals and, to aid in the prevention of contamination with disease causing organisms, the following proctices should be observed:

(a) Cleaned and disinfected containers should be used in collecting the eggs, and precautions taken to prevent contamination from organisms that may be present on the hands or clothing of the person making the collection.

(b) Dirty eggs should not be used for hatching purposes and should be collected in a separate container from hatching eggs. Slightly soiled eggs may be dry cleaned by hand or motor driven buffer.

(c) The visibly clean eggs should be fumigated as described in §147.25(a) as soon as possible after collection. The fumigated eggs shall be stored in a cool place at temperatures which will prevent the eggs from sweating at any time.

(d) Egg handlers should thoroughly wash their hands with soap and water and change to clean outer garments prior to handling the sanitized eggs. Sanitized eggs should be immediately removed from the cleaning and grading area and preferably removed to a separate clean and sanitized room. A wall-installed fumigation cabinet (or authorized sanitizing equipment) through which eggs can be passed from the receiving and cleaning area to the sanitary packing and storage areas is a good practice.

(e) The sanitized eggs should be placed in new flats or sanitized reusable flats or racks. New or clean, fumigated, used cases should be utilized for packing. Proper temperature and humidity in the egg cleaning, packing, and storage areas should be maintained. Eggs should be stored no longer than necessary before setting.

(f) The entire egg processing area should be cleaned and sanitized daily on a routine basis; dust, insects, feathers, and other airborn debris should be effectively controlled to prevent recontamination of sanitized eggs. Ink stamps and pad shall be maintained in a sterile condition.

(g) The egg processing building or area should be designed, located, and constructed of such materials as to assure that proper egg sanitation procedures can be carried out, and that the building itself can be easily, effectively, and routinely sanitized. The egg processing building or area should be considered part of a hatchery and the same construction details and physical and personnel sanitation requirements implemented.

§147.23 Hatchery sanitation.

An effective program for the prevention and control of salmonella and other infections should include the following measures:

(a) The hatchery building should be arranged so that separate rooms, with separate ventilation, are provided for each of the four operations: Egg receiving, incubation and hatching, holding of baby poultry, and disposal of offal and cleaning of trays. These rooms should be placed under isolation so that admission is granted only to specifically authorized personnel who have taken proper precautions to prevent introduction of disease.

(b) The hatchery rooms, and tables, racks, and other equipment in them should be thoroughly cleaned and disinfected frequently. All hatchery wastes and offal should be burned or otherwise properly disposed of, and the containers used to remove such materials should be cleaned and sterilized after each use.

(c) The hatching compartments of incubators, including the hatching trays, should be thoroughly cleaned and fumigated after each hatch.

(d) Only clean eggs should be used for hatching purposes. All eggs set should be fumigated prior to setting or as soon as possible (preferably within 12 hours) after they are placed in the incubator. They should also be fumigated after transfer to a separate hatcher. (See §147.25(d).)

(e) Only new or clean, fumigated egg cases should be used for transportation of hatching eggs. Soiled egg case fillers should be destroyed.

(f) Day-old chicks, poults, or other newly hatched poultry should be distributed in clean, new boxes. All crates and vehicles used for transporting started or adult birds should be cleaned and disinfected after each use.

§147.24 Cleaning and disinfecting.

The following procedures are recommended:

(a) In the poultry houses, hatchery rooms, and delivery trucks:

(a)(1) Settle dust by spraying lightly with the disinfectant to be used.

(a)(2) Remove all litter and droppings to an isolated area where there is no opportunity for dissemination of any infectious disease organisms that may be present.

(a)(3) Scrub the walls, floors, and equipment with a hot soapy water solution. Rinse to remove soap.

(a)(4) Spray with a disinfectant which is registered by the Environmental Protection Agency as germicidal, fungicidal, pseudomonocidal, and tuberculocidal, in accordance with the specifications for use, as shown on the label of such disinfectant.

(b) In the hatchers:

(b)(1) Remove trays and all controls and fans for separate cleaning. The ceiling, walls, and floors should be thoroughly wetted with a stream of water; then scubbed with a hard bristle brush. Rinse until there is no longer any deposit on the walls, particularly near the fan opening.

(b)(2) Replace the cleaned fans and controls. Replace the trays, preferably still wet from cleaning, and bring the incubator to normal operating temperature.

(b)(3) The hatcher should be fumigated as described in §147.25(e) prior to the transfer of the eggs.

(c) If the same machine is used for incubating and hatching, the entire machine should be cleaned after each hatch. A vacuum cleaner should be used to remove dust and down from the egg trays; then the entire machine should be vacuumed, mopped, and fumigated according to the procedures described in §147.25(b) (3), (4), and (5).

§147.25 Fumigation.

Fumigation is recommended for sanitizing eggs and hatchery equipment as an essential part of a sanitation program.

(a) Fumigation of clean eggs after collection should be done as follows:

(a)(1) Provide a room or cabinet proportionate to the number of eggs to be handled. The room or cabinet should be airtight and must be equipped with a fan to circulate the gas during fumigation and to expel it after fumigation.

(a)(2) The eggs should be placed on wire racks, in wire baskets, or on cup-type egg flats stacked outside of the egg cases (to permit air circulation) and exposed to circulating formaldehyde gas.

(a)(3) Formaldehyde gas is provided by mixing 0.6 gram of potassium permanganate with 1.2 cc of formalin (37.5 percent) for each cubic foot of space in the room. The ingredients should be mixed in an earthenware or enamelware container having a capacity at least 10 times the volume of the total ingredients.

(a)(4) Circulate the gas within the room for 20 minutes; then expel.

(a)(5) The temperature in the cabinet during fumigation should be at least 70°F., and the relative humidity above 70 percent.

(b) Eggs should be fumigated at the hatchery prior to setting or as soon as possible after setting (preferably within 12 hours). Single or repeated fumigation of eggs in the setter may be practiced, but the fumigation schedule should be such that no eggs are fumigated during the period from the 24th to the 84th hour of incubation. The following procedure should be used:

(b)(1) Determine the size of the incubator by multiplying the length times the width times the height.

(b)(2) After setting the eggs and allowing temperature and humidity to regain normal operating levels, release formaldehyde gas into the incubator.

(b)(3) For each cubic foot of space in the incubator, use 0.4 grams of potassium permanganate and 0.8 cc of formalin (37.5 percent). For mixing the fumigant, use an earthenware or enamelware container having the capacity of at least 10 times the volume of the total ingredients.

(b)(4) Close vents and doors but keep circulating fan operating, and continue fumigation for 20 minutes with normal operating temperature and humidity.

(b)(5) After 20 minutes of fumigation, the vents should be opened to the normal operating positions to release the gas.

(c) Eggs which have not been fumigated in the hatchery as described in paragraph (b) of this section should be fumigated after the 84th hour of incubation. The procedure described in paragraph (b) of this section should be followed.

(d) All eggs should be fumigated after transfer to a separate hatcher, preferably as soon as the temperature and humidity regain normal operating levels. The procedure described in paragraph (b) of this section should be followed.

(e) Empty hatchers should be fumigated between each hatch. After the interior of the hatcher has been thoroughly cleaned and the cleaned trays returned, the following procedure should be followed:

(e)(1) After temperature and humidity are brought to normal operating levels, use 0.6 grams of potassium permanganate and 1.2 cc of formalin (37.5 percent) per cubic foot of space in the hatcher.

(e)(2) Close the doors and vents and leave closed at least 3 hours, preferably overnight.

(f) The cheesecloth method of fumigation described in this paragraph may be used in lieu of the chemical method described in paragraph (b) of this section, using 0.6 cc of formalin (37.5 percent) per cubic foot of space in the incubator, or in lieu of the chemical method described in paragraph (e) of this section, using 0.9 cc of formalin (37.5 percent) for each cubic foot of space in the empty hatcher.

(f)(1) Enough cheesecloth should be used to absorb all of the formalin that is to be used for the fumigation.

(f)(2) The formalin-saturated cheesecloth should be hung in the cabinet in such a manner as to permit the circulating air to evaporate all the formalin. This will require longer than 20 minutes.

(f)(3) Care should be taken to prevent the cheesecloth from blocking the air movement created by the fans.

(f)(4) The cheesecloth method is not suitable for still air machines.

§147.26 Procedures for establishing isolation and maintaining sanitation and good management practices for the control of salmonella and mycoplasma infections.

(a) The following procedures are required for participation under the U.S. Sanitation Monitored, U.S. M. Gallisepticum Clean, and U.S. M. Synoviae Clean classifications:

(a)(1) Allow no visitors except under controlled conditions which insure sanitation. Such conditions shall be approved by the Official State Agency and the Service;

(a)(2) Maintain breeder flocks on farms free from market birds, or follow proper isolation procedures as approved by the Official State Agency;

(a)(3) Elimiate other domesticated fowl from breeder farm;

(a)(4) Dispose of all dead birds by burning, deep burial, or by putting them into special disposal pits.

(b) Recommended procedures:

(b)(1) Avoid the introduction of salmonella, Mycoplasma gallisepticum, or Mycoplasma synoviae infected poultry;

(b)(2) Prevent indirect transmission from outside sources through contaminated equipment, footwear, clothing, vehicles, or other mechanical means;

(b)(3) Provide adequate isolation of breeder flocks to avoid airborn transmission from infected flocks;

(b)(4) Minimize contact of breeder flocks with free-flying birds;

(b)(5) Keep the rodent population and other pests under control;

(b)(6) Tailor vaccination programs to needs of farm and area;

(b)(7) Clean and disinfect equipment after each use;

(b)(8) Provide clean footwear and provide an adequate security program;

(b)(9) Clean and disinfect houses before introducing a new flock;

(b)(10) Use well-drained range;

(b)(11) Use clean, dry litter free of mold;

(b)(12) Keep accurate records of death losses;

(b)(13) Seek services of veterinary diagnostician if unaccountable mortality or signs of disease occur;

(b)(14) Adopt and maintain a clean-egg program;

(b)(15) Use only crates and vehicles that have been cleaned and disinfected in accordance with the provisions of §147.24(a) to haul live poultry to and from the premises.

§147.27 Procedures recommended to prevent the spread of diseases by artificial insemination of turkeys.

(a) The vehicle transporting the insemination crew should be left as far as practical from the turkey pens.

(b) The personnel of the insemination crew should observe personal cleanliness, including the following sanitary procedures:

(b)(1) Outer clothing should be changed between visits to different premises so that clean clothing is worn upon entering each premises. The used apparel should be kept separate until laundered. This also applies to gloves worn while handling turkeys;

(b)(2) Boots or footwear should be cleaned and disinfected between visits to different premises;

(b)(3) Disposable caps should be provided and discarded after use on each premises.

(c) The use of individual straw or similar technique is highly recommended. Insemination equipment which is to be reused should be cleaned and disinfected before reusing. Equipment used for the convenience of the workers should not be moved from premises to premises.

(d) No obviously diseased flock should be inseminated. If evidence of active disease is noted after insemination is begun, operations should be stopped and the hatchery notified.

(f) Care should be taken during the collection of semen to prevent fecal contamination. If fecal material is present, it should be removed before the semen is collected. Likewise, care should be taken not to introduce fecal material into the oviduct of the hen.

Subpart D - [Reserved]

Subpart E - Procedure For Changing National Poultry Improvement Plan

§147.41 Definitions.

Except where the context otherwise requires, for the purposes of this subpart the following terms shall be construed, respectively, to mean:

(a) Plan or NPIP. The National Poultry Improvement Plan.

(b) Plan Conference. A meeting convened for the purpose of recommending changes in the provisions of the Plan.

(c) Department. The U.S. Department of Agriculture.

(d) Service. The Animal and Plant Health Inspection Service-Veterinary Services of the Department.

(e) State. Any State, the District of Columbia, or Puerto Rico.

(f) Egg-type chickens. Chickens bred for the primary purpose of producing eggs for human consumption.

(g) Meat-type chickens. Chickens bred for the primary purpose of producing meat.

(h) Waterfowl. Domesticated fowl that normally swim, such as ducks and geese.

(i) Exhibition Poultry. Domesticated fowl which are bred for the combined purposes of meat or egg production and competitive showing.

(j) Game birds. Domesticated fowl, such as pheasant, partridge, quail, grouse, and guineas, but not doves and pigeons.

§147.42 General.

Changes in this subchapter shall be made in accordance with the procedure described in this subpart: Provided, That the Department reserves the right to make changes in this subchapter without observance of such procedure when such action is deemed necessary in the public interest.

§147.43 General Conference Committee.

(a) The General Conference Committee shall consist of the Assistant Secretary of Agriculture for Marketing and Transportation Services, or his/her designee, one member-at-large who is a participant in the National Poultry Improvement Plan and who shall be designated as vice chairperson and one member to be elected, as provided in paragraph (b) of this section, from each of the following regions:

(a)(1) North Atlantic: Maine, New Hampshire, Vermont, Massachusetts, Rhode Island, Connecticut, New York, New Jersey, and Pennsylvania.

(a)(2) East North Central: Ohio, Indiana, Illinois, Michigan, and Wisconsin.

(a)(3) West North Central: Minnesota, Iowa, Missouri, North Dakota, South Dakota, Nebraska, and Kansas

(a)(4) South Atlantic: Delaware, District of Columbia, Maryland, Virginia, West Virginia, North Carolina, South Carolina, Georgia, Florida, and Puerto Rico.

(a)(5) South Central: Kentucky, Tennessee, Alabama, Mississippi, Arkansas, Louisiana, Oklahoma, and Texas.

(a)(6) Western: Montana, Idaho, Wyoming, Colorado, New Mexico, Arizona, Utah, Nevada, Washington, Oregon, California, Alaska, and Hawaii.

(b) The regional Committee members and their alternates will be elected by the official delegates of the respective regions and the member-at-large will be elected by all official delegates. There shall be at least two nominees for each position, and the voting shall be by secret ballot.

(c) Three regional members shall be elected at each Plan Conference. All members shall serve for a period of four years, subject to the continuation of the Committee by the Secretary of Agriculture, and may not succeed themselves. When there is a vacancy for the member-at-large position, the General Conference Committee shall make an interim appointment and the appointee shall serve until the next Plan Conference at which time an election will be held.

(d) The duties and functions of the General Conferrence Committee shall be as follows:

(d)(1) Assist the Department in planning, organizing, and conducting the biennial National Poultry Improvement Plan Conference.

(d)(2) Recommend whether new proposals (i.e., proposals that have not been submitted as provided in §147.44) should be considered by the delegates to the Plan Conference.

(d)(3) During the interim between Plan Conferences, represent the cooperating State in:

(d)(3)(i) Advising the Department with respect to administrative procedures and interpretations of the Plan provisions as contained in 9 CFR.

(d)(3)(ii) Assisting the Department in evaluating comments received from interested person concerning proposed amendments to the Plan provisions.

(d)(3)(iii) Recommending to the Secretary of Agriculture any changes in the provisions of the Plan as may be necessitated by unforeseen conditions when postponement until the next Plan Conference would seriously impair the operation of the program. Such recommendations shall remain in effect only until confired or rejected by the next Plan Conference, or until rescinded by the committee.

(d)(4) Serve as a forum for the study of problems relating to poultry health and as the need arises, to make specific recommendations to the Secretary of Agriculture concerning ways in which the Department may assist the industry in solving these problems. (Sec. 101(b), Pub. L. 425, 78th Cong. 58 Stat. 734, as amended 7 U.S.C. 429)

§147.44 Submitting, compiling, and distributing proposed changes.

(a) Changes in this subchapter may be proposed by any participant, Official State Agency, the Department, or other interested person or industry organization.

(b) Except as provided in §147.43(d)(1), proposed changes shall be submitted in writing so as to reach the Service not later than 150 days prior to the opening date of the Plan Conference, and participants in the Plan shall submit their proposed changes through their Official State Agency.

(c) The name of the proponent shall be indicated in each proposed change when submitted. Each proposal should be accompained by a brief supporting statement.

(d) The Service will notify all persons on the NPIP mailing lists concerning the dates and general procedure of the conference. Hatchery and dealer participants will be reminded of their privilege to submit proposed changes and to request copies of all the published proposed changes.

(e) The proposed changes, together with the names of the proponents and supporting statements, will be compiled by the Service and issued in processed form. When two or more similar changes are submitted, the Service will endeavor to unify them into one proposal acceptable to each proponent. Copies will be distributed to officials of the Official State Agencies cooperating in the NPIP. Additional copies will be made available for meeting individual requests.

§147.45 Official delegates.

Each cooperating State shall be entitled to one official delegate for each of the programs prescribed in Subparts B, C, D, and E of Part 145 of this chapter in which it has one or more participants at the time of the Conference. The official delegates shall be elected by a representative group of participating industry members and be certified by the Official State Agency. It is recommended but not required that the official delegates be Plan participants. Each official delegate shall endeavor to obtain, prior to the Conference, the recommendations of industry members of his State with respect to each proposed change.

§147.46 Committee consideration of proposed changes.

(a) The following four committees shall be established to give preliminary consideration to the proposed changes falling in their respective fields:

(a)(1) Egg-type chickens.

(a)(2) Meat-type chickens.

(a)(3) Turkeys.

(a)(4) Waterfowl, exhibition poultry, and game birds.

(b) Each official delegate shall be appointed a voting member in one of the committees specified in paragraph (a) of this section.

(c) Since several of the proposals may be interrelated, the committees shall consider them as they may relate to others, and feel free to discuss related proposals with other committees.

(d) The committees shall make recommendations to the conference as a whole concerning each proposal. The committee report shall show any proposed change in wording and the record of the vote on each proposal, and suggest an effective date for each proposal recommended for adoption. The individual committee reports shall be submitted to the chairman of the conference, who will combine them into one report showing, in numerical sequence, the committee recommendations on each proposal.

(e) The committee meetings shall be open to any interested person. Advocates for or against any proposal should feel free to appear before the appropriate committee and present their views.

§147.47 Conference consideration of proposed changes.

(a) The chairman of the conference shall be a representative of the Department.

(b) At the time designated for voting on proposed changes by the official delegates, the chairman of the General Conference Committee and the four committee chairmen shall sit at the speaker's table and assist the chairman of the conference.

(c) Each committee chairman shall present the proposals which his committee approves or recommends for adoption as follows: "Mr. Chairman, the committee for egg-type chickens recommends the adoption of Proposal No. ____, for the following reasons (stating the reasons): I move the adoption of Proposal No. _____." A second will then be called for. If the recommendation is seconded, discussion and a formal vote will follow.

(d) Each committee chairman shall present the proposals which his committee does not approve as follows: "Mr. Chairman, the Committee for egg-type chickens does not approve Proposal No. ____." The chairman will then ask if any official delegate wishes to move for the adoption of the proposal. If moved and seconded, the proposal is subject to discussion and voted. If there is no motion for approval, or if moved but not seconded, there can be no discussion or vote.

(e) Discussion on any motion must be withheld until the motion has been properly seconded, except that the delegate making the motion is privileged, if he desires, to give reasons for his motion at the time of making it. To gain the floor for a motion or for discussion on a motion, the official delegate in the case of a motion, or anyone in case of discussion on a motion, shall rise, address the chair, give his name and State, and be recognized by the chair before proceeding further. While it is proper to accept motions only from official delegates and to limit voting only to such delegates, it is, however, equally proper to accept discussion from anyone interested. To conserve time, discussion should be pointed and limited to the pertinent features of the motion.

(f) Proposals that have not been submitted in accordance with §147.44 will be considered by the conference only with the unanimous consent of the General Conference Committee. Any such proposals must be referred to the appropriate committee for consideration before being presented for acton by the conference.

(g) Voting will be by States, and each official delegate, as determined by §147.45, will be allowed one vote on each proposal pertaining to the program prescribed by the subpart which he represents.

(h) A rollcall of States for a recorded vote will be used when requested by a delegate or at the discretion of the chairman.

(i) All motions on proposed changes shall be for adoption.

(j) Proposed changes shall be adopted by a majority vote of the official delegates present and voting.

(k) The conference shall be open to any interested person.

§147.48 Approval of conference recommendations by the Department.

Proposals adopted by the official delegates will be recommended to the Department for incorporation into the provisions of the NPIP. The Department reserves the right to approve or disapprove the recommendations of the conference as an integral part of its sponsorship of the National Poultry Improvement Plan.

NOTE: The recordkeeping and/or reporting requirements contained herein have been approved by the Office of Management and Budget in accordance with the Federal Reports Act of 1942.

United States
Department of Agriculture

National Poultry Improvement Plan
BARC-East, Bldg. 265
Beltsville, Md. 20705

OFFICIAL BUSINESS
Penalty for Private Use, $300

Postage and Fees Paid
U.S. Department of Agriculture
AGR-101